

Dobrushin Conditions for Systematic Scan with Block Dynamics*

Kasper Pedersen
 Department of Computer Science
 University of Liverpool
 Liverpool L69 3BX, UK
 k.pedersen@csc.liv.ac.uk

February 2, 2008

Abstract

We study the mixing time of systematic scan Markov chains on finite spin systems. It is known that, in a single site setting, the mixing time of systematic scan can be bounded in terms of the influences sites have on each other. We generalise this technique for bounding the mixing time of systematic scan to block dynamics, a setting in which a (constant size) set of sites are updated simultaneously. In particular we consider the parameter α , corresponding to the maximum influence on any site, and show that if $\alpha < 1$ then the corresponding systematic scan Markov chain mixes rapidly. As applications of this method we prove $O(\log n)$ mixing of systematic scan (for any scan order) for heat-bath updates of edges for proper q -colourings of a general graph with maximum vertex-degree Δ when $q \geq 2\Delta$. We also apply the method to improve the number of colours required in order to obtain mixing in $O(\log n)$ scans for systematic scan for heat-bath updates on trees, using some suitable block updates.

1 Introduction

This paper is concerned with the study of finite *spin systems*. A spin system is composed of a set of sites and a set of spins, both of which will be finite throughout this paper. The interconnection between the sites is determined by an underlying graph. A configuration of the spin system is an assignment of a spin to each site. If there are n sites and q available spins then this gives rise to q^n configurations of the system, however some configurations may be illegal. The specification of the system determines how the spins interact with each other at a local level, such that different local configurations on a subset of the graph may have different relative likelihoods. This interaction hence specifies a probability distribution, π , on the set of configurations. One class of configurations that receive much attention in theoretical computer science is *proper q -colourings* of graphs. A proper colouring is a configuration where no two adjacent sites are assigned the same colour. One important example of a spin system is when the set of legal configurations is the set of all proper q -colourings of the underlying graph and π is the uniform distribution on this set. In statistical physics the spin system corresponding to proper q -colourings is known as the q -state anti-ferromagnetic Potts model at zero temperature.

*This work was partly funded by EPSRC projects GR/T07343/02 and GR/S76168/01.

Sampling from π is a computationally challenging task. It is, however, an important one and is often carried out by simulating some suitable random *dynamics* on the set of configurations. Such a dynamics must have the following two properties

1. the dynamics eventually converges to π , and
2. the rate of convergence (*mixing time*) is polynomial in the number of sites.

It is generally straightforward to ensure that a dynamics converges to π but much harder provide good upper bounds on the rate of convergence, which is what we will be concerned with in this paper.

Arguably the simplest dynamics is the heat-bath Glauber dynamics which, at each step, selects a site uniformly at random and updates the spin assigned to that site by drawing a new spin from the distribution on the spin of the selected site induced by π . This procedure is repeated until the distribution of the Markov chain is sufficiently close to π using some suitable measure of closeness between probability distributions. This dynamics falls under a family of Markov chains that we call *random update Markov chains*. We say that a Markov chain is a random update Markov chain if the sites are updated in a random order. This type of Markov chain has been frequently studied in theoretical computer science and much is known about the mixing time of various random update Markov chains.

An alternative to random update Markov chains is to construct a Markov chain that cycles through (and updates the spin according to the induced distribution) the sites (or subsets of sites) in a deterministic order. We call this a *systematic scan Markov chain* (or systematic scan for short). Although systematic scan updates the sites in a deterministic order it remains a random process since the procedure used to update the spin assigned to a site is randomised, as specified by the appropriate induced distribution. Systematic scan may be more intuitively appealing than random update in terms of implementation, however until recently little was known about the convergence rates of this type of dynamics. It remains important to know how many steps one needs to simulate a systematic scan for in order for it to become sufficiently close to its stationary distribution and recently there has been an interest among computer scientists in investigating various approaches for analysing the mixing time of systematic scan Markov chains, see e.g. Dyer, Goldberg and Jerrum [5, 7] and Bordewich, Dyer and Karpinski [2]. In this paper we present a new method for analysing the mixing time of systematic scan Markov chains, which is applicable to any spin system. As applications of this method we improve the known parameters required for rapid mixing of systematic scan on

1. proper colourings of general graphs and
2. proper colourings of trees.

A key ingredient in our method for proving mixing of systematic scan is to work with a *block dynamics*. A block dynamics is a dynamics in which we allow a set of sites to be updated simultaneously as opposed to updating one site at a time as in the description of the Glauber dynamics above. Block dynamics is not a new concept and it was used in the mid 1980s by Dobrushin and Shlosman [4] in their study of conditions that imply uniqueness of the Gibbs measure of a spin system, a topic closely related to studying the mixing time of Markov chains (see for example Weitz's PhD thesis [16]). More recently, a block dynamics has been used by Weitz [17] when, in a generalisation of the

work of Dobrushin and Shlosman, studying the relationship between various *influence parameters* (also in the context of Gibbs measures) within spin systems and using the influence parameters to establish conditions that imply mixing. Using an influence parameter to establish a condition which implies mixing of systematic scan is a key aspect of the method presented in this paper as we will discuss below. Dyer, Sinclair, Vigoda and Weitz [8] have also used a block dynamics in the context of analysing the mixing time of a Markov chain for proper colourings of the square lattice. Both of these papers consider a random update Markov chain, however several of ideas and techniques carry over to systematic scan as we shall see.

We will bound the mixing time of systematic scan by studying the *influence* that the sites of the graph have on each other. This technique is well-known and the influence parameters generalised by Weitz [17]: “the influence *on* a site is small” (originally attributed to Dobrushin [3]) and “the influence *of* a site is small” (originally Dobrushin and Shlosman [4]) both imply mixing of the corresponding random update Markov chain. It is worth pointing out that a condition of the form “if the influence *on* a site is small then the corresponding dynamics converges to π quickly” is known as a Dobrushin condition. In the context of systematic scan, Dyer et al. [5] point out that, in a single site setting, the condition “the influence *on* a site is small” implies rapid mixing of systematic scan. Our method for proving rapid mixing of systematic scan is a generalisation of this influence parameter to block dynamics.

We now formalise the concepts above and state our results. Let $C = \{1, \dots, q\}$ be the set of spins and $G = (V, E)$ be the underlying graph of the spin system where $V = \{1, \dots, n\}$ is the set of sites. We associate with each site $i \in V$ a positive weight w_i . Let Ω^+ be the set of all configurations of the spin system and $\Omega \subseteq \Omega^+$ be the set of all legal configurations. Then let π be a probability distribution on Ω^+ whose support is Ω i.e., $\{x \in \Omega^+ \mid \pi(x) > 0\} = \Omega$. If $x \in \Omega^+$ is a configuration and $j \in V$ is a site then x_j denotes the spin assigned to site j in configuration x . For each site $j \in V$, let S_j denote the set of pairs $(x, y) \in \Omega^+ \times \Omega^+$ of configurations that only differ on the spin assigned to site j , that is $x_i = y_i$ for all $i \neq j$.

We will use Weitz’s [17] notation for block dynamics, although we only consider a finite collection of blocks. Define a collection of m blocks $\Theta = \{\Theta_k\}_{k=1, \dots, m}$ such that each block $\Theta_k \subseteq V$ and Θ covers V , where we say that Θ covers V if $\bigcup_{k=1}^m \Theta_k = V$. One site may be contained in several blocks and the size of each block is not required to be the same, we do however require that the size of each block is bounded independently of n . For any block Θ_k and a pair of configurations $x, y \in \Omega^+$ we write “ $x = y$ on Θ_k ” if $x_i = y_i$ for each $i \in \Theta_k$ and similarly “ $x = y$ off Θ_k ” if $x_i = y_i$ for each $i \in V \setminus \Theta_k$. We also let $\partial\Theta_k = \{i \in V \setminus \Theta_k \mid \exists j \in \Theta_k : \{i, j\} \in E(G)\}$ denote the set of sites adjacent to but not included in Θ_k ; we will refer to $\partial\Theta_k$ as the *boundary* of Θ_k .

With each block Θ_k , we associate a transition matrix $P^{[k]}$ on state space Ω^+ satisfying the following two properties:

1. If $P^{[k]}(x, y) > 0$ then $x = y$ off Θ_k , and also
2. π is invariant with respect to $P^{[k]}$.

Property 1 ensures that an application of $P^{[k]}$ moves the state of the system from from one configuration to another by only updating the sites contained in the block Θ_k and Property 2 ensures that any dynamics composed solely of transitions defined by $P^{[k]}$ converges to π . While the requirements of Property 1 are clear we take a moment to discuss what we mean in Property 2. Consider the following two step process in which some configuration x is initially drawn from π and then a configuration y is drawn from $P^{[k]}(x)$ where $P^{[k]}(x)$ is the distribution on configurations resulting from

applying $P^{[k]}$ to a configuration x . We then say that π is invariant with respect to $P^{[k]}$ if for each configuration $\sigma \in \Omega^+$ we have $\Pr(x = \sigma) = \Pr(y = \sigma)$. That is the distribution on configurations generated by the two-step process is the same as if only the first step was executed. In terms of our dynamics this means that once the distribution of the dynamics reaches π , π will continue be the distribution of the dynamics even after applying $P^{[k]}$ to the state of the dynamics. Our main result (Theorem 2) holds for any choice of update rule $P^{[k]}$ provided that it satisfies these two properties.

The distribution $P^{[k]}(x)$, which specifies how the dynamics updates block Θ_k , clearly depends on the specific update rule implemented as $P^{[k]}$. In order to make this idea more clear we describe one particular update rule, known as the *heat-bath* update rule. This example serves a dual purpose as it is a simple way to implement $P^{[k]}$ and we will make use of heat-bath updates in Sections 3 and 4 when applying our condition to specific spin systems. A heat-bath move on a block Θ_k given a configuration x is performed by drawing a new configuration from the distribution induced by π and consistent with the assignment of spins on the boundary of Θ_k . The two properties of $P^{[k]}$ hold for heat-bath updates since (1) only the assignment of the spin to the sites in Θ_k are changed and (2) the new configuration is drawn from an appropriate distribution induced by π . If the spin system corresponds to proper colourings of graphs then the distribution used in a heat-bath move is the uniform distribution the set of configurations that agree with x off Θ_k and where no edge containing a site in Θ_k is monochromatic.

With these definitions in mind we are ready to formally define a systematic scan Markov chain.

Definition 1. We let $\mathcal{M}_\rightarrow$ be a systematic scan Markov chain with state space Ω^+ and transition matrix $P_\rightarrow = \prod_{k=1}^m P^{[k]}$.

The stationary distribution of $\mathcal{M}_\rightarrow$ is π as discussed above, and it is worth pointing out that the definition of $\mathcal{M}_\rightarrow$ holds for *any* order on the set of blocks. We will refer to one application of $P_\rightarrow$ (that is updating each block once) as one *scan* of $\mathcal{M}_\rightarrow$. One scan takes $\sum_k |\Theta_k|$ *updates* and it is generally straight forward to ensure, via the construction of the set of blocks, that this sum is of order $O(n)$.

We will be concerned with analysing the mixing time of systematic scan Markov chains, and consider the case when $\mathcal{M}_\rightarrow$ is ergodic. Let $\mathcal{M}$ be any ergodic Markov chain with state space Ω^+ and transition matrix P . By classical theory (see e.g. Aldous [1]) $\mathcal{M}$ has a unique stationary distribution, which we will denote π . The mixing time from an initial configuration $x \in \Omega^+$ is the number of steps, that is applications of P , required for $\mathcal{M}$ to become sufficiently close to π . Formally the mixing time of $\mathcal{M}$ from an initial configuration $x \in \Omega^+$ is defined, as a function of the deviation ε from stationarity, by

$$\text{Mix}_x(\mathcal{M}, \varepsilon) = \min\{t > 0 : d_{\text{TV}}(P^t(x, \cdot), \pi(\cdot)) \leq \varepsilon\}$$

where

$$d_{\text{TV}}(\theta_1, \theta_2) = \frac{1}{2} \sum_i |\theta_1(i) - \theta_2(i)| = \max_{A \subseteq \Omega^+} |\theta_1(A) - \theta_2(A)|$$

is the total variation distance between two distributions θ_1 and θ_2 on Ω^+ . The mixing time $\text{Mix}(\mathcal{M}, \varepsilon)$ of $\mathcal{M}$ is then obtained by maximising over all possible initial configurations

$$\text{Mix}(\mathcal{M}, \varepsilon) = \max_{x \in \Omega^+} \text{Mix}_x(\mathcal{M}, \varepsilon).$$

We say that $\mathcal{M}$ is *rapidly mixing* if the mixing time of $\mathcal{M}$ is polynomial in n and $\log(\varepsilon^{-1})$.

We will now formalise the notion of “the influence *on* a site” in order to state our condition for rapid mixing of systematic scan. For any pair of configurations (x, y) let $\Psi_k(x, y)$ be a coupling of the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$ which we will refer to as “updating block Θ_k ”. Recall that a coupling $\Psi_k(x, y)$ of $P^{[k]}(x)$ and $P^{[k]}(y)$ is a joint distribution on $\Omega^+ \times \Omega^+$ whose marginal distributions are $P^{[k]}(x)$ and $P^{[k]}(y)$. That is

$$\forall \sigma \in \Omega^+ \quad \Pr_{x' \in P^{[k]}(x)}(x' = \sigma) = \sum_{\tau \in \Omega^+} \Pr_{(x', y') \in \Psi_k(x, y)}(x' = \sigma, y' = \tau)$$

and

$$\forall \tau \in \Omega^+ \quad \Pr_{y' \in P^{[k]}(y)}(y' = \tau) = \sum_{\sigma \in \Omega^+} \Pr_{(x', y') \in \Psi_k(x, y)}(x' = \sigma, y' = \tau)$$

where we write $(x', y') \in \Psi_k(x, y)$ when the pair of configurations (x', y') is drawn from $\Psi_k(x, y)$. Weitz in [17] states his conditions for general metrics whereas we will use Hamming distance, which is also how the corresponding condition is defined in Dyer et al. [5]. This choice of metric allows us to define the influence of a site i on a site j under a block Θ_k , which we will denote $\rho_{i,j}^k$, as the maximum probability that two coupled Markov chains differ at the spin of site j following an update of Θ_k starting from two configurations that only differ at the spin on site i . That is

$$\rho_{i,j}^k = \max_{(x, y) \in S_i} \{\Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j)\}.$$

Then let α be the total (weighted) influence *on* any site in the graph site defined by

$$\alpha = \max_k \max_{j \in \Theta_k} \sum_i \frac{w_i}{w_j} \rho_{i,j}^k.$$

We point out that our definition of $\rho_{i,j}^k$ is not the standard definition of ρ used in the literature (see for example Simon [14] or Dyer et al. [5]) since the coupling $\Psi_k(x, y)$ is explicitly included. In the block setting it is, however, necessary to include the coupling directly in the definition of ρ as we will discuss in Section 5. In Section 5 we also show that the condition $\alpha < 1$ is a generalisation of the corresponding condition in Dyer et al. [5] in the sense that if each block contains exactly one site and the coupling minimises the Hamming distance then the conditions coincide. Our main theorem, which is proved in Section 2, states that if the influence on a site is sufficiently small then the systematic scan Markov chain $\mathcal{M}_\rightarrow$ mixes in $O(\log n)$ scans.

Theorem 2. *Suppose $\alpha < 1$. Then*

$$\text{Mix}(\mathcal{M}_\rightarrow, \varepsilon) \leq \frac{\log(n\varepsilon^{-1})}{1 - \alpha}.$$

As previously stated we will apply Theorem 2 to two spin systems corresponding to proper q -colourings of graphs in order to improve the parameters for which systematic scan mixes. In both applications we restrict the state space of the Markov chains to the set of proper colourings, Ω , of the underlying graph. Firstly we allow the underlying graph to be any finite graph with maximum vertex-degree Δ . Previously, the least number of colours for which systematic scan was known to mix in $O(\log n)$ scans was $q > 2\Delta$ and when $q = 2\Delta$ the best known bound on the mixing time was $O(n^2 \log n)$ scans due to Dyer et al. [5]. For completeness we pause to mention that the least number of colours required for rapid mixing of a random update Markov chain is $q > 11/6\Delta$ due

Table 1: Optimising the number of colours using blocks

Δ	h	ξ	$f(\Delta)$	$\lceil \Delta + 2\sqrt{\Delta - 1} \rceil$
3	15	$\frac{4}{7}$	5	6
4	3	$\frac{5}{11}$	7	8
5	12	$\frac{5}{11}$	8	9
6	3	$\frac{1}{2}$	10	11
7	7	$\frac{10}{23}$	11	12
8	13	$\frac{1}{3}$	12	14
9	85	$\frac{5}{19}$	13	15
10	5	$\frac{5}{19}$	15	16

to Vigoda [15]. In Section 3 we consider the following Markov chain, *edge scan* denoted $\mathcal{M}_{\text{edge}}$, updating the endpoints of an edge during each update. Let $\Theta = \{\Theta_k\}_{k=1,\dots,m}$ be a set of edges in G such that Θ covers V . Using the above notation, $P^{[k]}$ is the transition matrix for performing a heat-bath move on the endpoints of the edge Θ_k and the transition matrix of $\mathcal{M}_{\text{edge}}$ is $\Pi_{k=1}^m P^{[k]}$. We prove the following theorem, which improves the mixing time of systematic scan by a factor of n^2 for proper colourings of general graphs when $q = 2\Delta$ and matches the existing bound when $q > 2\Delta$.

Theorem 3. *Let G be a graph with maximum vertex-degree Δ . If $q \geq 2\Delta$ then*

$$\text{Mix}(\mathcal{M}_{\text{edge}}, \varepsilon) \leq \Delta^2 \log(n\varepsilon^{-1}).$$

Next, in Section 4, we restrict the class of graphs to trees. It is known that single site systematic scan mixes in $O(\log n)$ scans when $q > \Delta + 2\sqrt{\Delta - 1}$ and in $O(n^2 \log n)$ scans when $q = \Delta + 2\sqrt{\Delta - 1}$ is an integer; see e.g. Hayes [11] or Dyer, Goldberg and Jerrum [6]. More generally it is known that systematic scan for proper colourings of bipartite graphs mixes in $O(\log n)$ scans when $q \geq 1.76\Delta$ as $\Delta \rightarrow \infty$ due to Bordewich et al. [2]. Again, for completeness, we mention that the mixing time of a random update Markov chain for proper colourings on a tree mixes in $O(n \log n)$ updates when $q \geq \Delta + 2$, a result due to Martinelli, Sinclair and Weitz [13], improving a similar result by Kenyon, Mossel and Peres [12]. We will use a block approach to improve the number of colours required for mixing of systematic scan on trees. We construct the following set of blocks where the height h of the blocks is defined in Table 1. Let a block Θ_k contain a site r along with all sites below r in the tree that are at most $h - 1$ edges away from r . The set of blocks Θ covers the sites of the tree and we construct Θ such that no block has height less than h . $P^{[k]}$ is the transition matrix for performing a heat-bath move on block Θ_k and the transition matrix of the Markov chain $\mathcal{M}_{\text{tree}}$ is $\Pi_{k=1}^m P^{[k]}$ where m is the number of blocks. We prove the following theorem.

Theorem 4. *Let G be a tree with maximum vertex-degree Δ . If $q \geq f(\Delta)$ where $f(\Delta)$ is specified in Table 1 for small Δ then*

$$\text{Mix}(\mathcal{M}_{\text{tree}}, \varepsilon) = O(\log(n\varepsilon^{-1})).$$

We conclude the paper with a discussion, in Section 5, of the influence parameter α and how it relates to the corresponding parameters for the “influence on a site” in Weitz [17] and Dyer et al. [5]. In particular we will show that the condition in Weitz [17] does not imply mixing of systematic scan and that the condition in Dyer et al. [5] is a special case of our condition from Theorem 2.

2 Bounding the Mixing Time of Systematic Scan

This section will contain the proof of Theorem 2. The proof follows the structure of the proof from the single-site setting in Dyer et al. [5], which follows Föllmer's [9] account of Dobrushin's proof presented in Simon's book [14].

We will make use of the following definitions. For any function $f : \Omega^+ \rightarrow \mathbb{R}_{\geq 0}$ let $\delta_i(f) = \max_{(x,y) \in S_i} |f(x) - f(y)|$ and $\Delta(f) = \sum_{i \in V} w_i \delta_i(f)$. Also for any transition matrix P define (Pf) as the function from Ω^+ to $\mathbb{R}_{\geq 0}$ given by $(Pf)(x) = \sum_{x'} P(x, x') f(x')$. Finally let $\mathbf{1}_{i \notin \Theta_k}$ be the function given by

$$\mathbf{1}_{i \notin \Theta_k} = \begin{cases} 1 & \text{if } i \notin \Theta_k \\ 0 & \text{otherwise.} \end{cases}$$

We can think of $\delta_i(f)$ as the deviation from constancy of f at site i and $\Delta(f)$ as the aggregated deviation from constancy of f . Now, Pf is a function where $(Pf)(x)$ gives the expected value of f after making a transition starting from x . Intuitively, if t transitions are sufficient for mixing then $P^t f$ is a very smooth function. An application of $P^{[k]}$ fixes the non-constancy of f at the sites within Θ_k although possibly at the cost of increasing the non-constancy at sites on the boundary of Θ_k . Our aim is then to show that one application of $P_{\rightarrow}$ will on aggregate make f smoother i.e., decrease $\Delta(f)$. We will establish the following lemma, which corresponds to Corollary 12 in Dyer et al. [5], from which Section 3.3 of [5] implies Theorem 2.

Lemma 5. *If $\alpha < 1$ then*

$$\Delta(P_{\rightarrow} f) \leq \alpha \Delta(f).$$

We begin by bounding the effect on f from one application of $P^{[k]}$. The following lemma is a block-move generalisation of Proposition V.1.7 from Simon [14] and Lemma 10 from Dyer et al. [5].

Lemma 6. $\delta_i(P^{[k]} f) \leq \mathbf{1}_{i \notin \Theta_k} \delta_i(f) + \sum_{j \in \Theta_k} \rho_{i,j}^k \delta_j(f)$

Proof. Take $\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [f(x')]$ to be the expected value of $f(x')$ when a pair of configurations (x', y') are drawn from $\Psi_k(x, y)$. Since $\Psi_k(x, y)$ is a coupling of the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$, the distribution $P^{[k]}(x)$ and the first component of $\Psi_k(x, y)$ are the same and hence

$$\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [f(x')] = \mathbf{E}_{x' \in P^{[k]}(x)} [f(x')] \quad (1)$$

and the same fact holds for the distribution $P^{[k]}(y)$ so

$$\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [f(y')] = \mathbf{E}_{y' \in P^{[k]}(y)} [f(y')]. \quad (2)$$

Using (1), (2) and linearity of expectation we have

$$\begin{aligned}
\delta_i(P^{[k]}f) &= \max_{(x,y) \in S_i} \left| (P^{[k]}f)(x) - (P^{[k]}f)(y) \right| \\
&= \max_{(x,y) \in S_i} \left| \sum_{x'} P^{[k]}(x, x') f(x') - \sum_{y'} P^{[k]}(y, y') f(y') \right| \\
&= \max_{(x,y) \in S_i} \left| \mathbf{E}_{x' \in P^{[k]}(x)} [f(x')] - \mathbf{E}_{y' \in P^{[k]}(y)} [f(y')] \right| \\
&= \max_{(x,y) \in S_i} \left| \mathbf{E}_{(x',y') \in \Psi_k(x,y)} [f(x')] - \mathbf{E}_{(x',y') \in \Psi_k(x,y)} [f(y')] \right| \\
&= \max_{(x,y) \in S_i} \left| \mathbf{E}_{(x',y') \in \Psi_k(x,y)} [f(x') - f(y')] \right| \\
&\leq \max_{(x,y) \in S_i} \mathbf{E}_{(x',y') \in \Psi_k(x,y)} [|f(x') - f(y')|] \\
&\leq \max_{(x,y) \in S_i} \mathbf{E}_{(x',y') \in \Psi_k(x,y)} \left[\sum_{j \in V} |f(x'_1 \dots x'_j y'_{j+1} \dots y'_n) - f(x'_1 \dots x'_{j-1} y'_j \dots y'_n)| \right] \\
&= \max_{(x,y) \in S_i} \sum_{j \in V} \mathbf{E}_{(x',y') \in \Psi_k(x,y)} [|f(x'_1 \dots x'_j y'_{j+1} \dots y'_n) - f(x'_1 \dots x'_{j-1} y'_j \dots y'_n)|].
\end{aligned}$$

Notice that $x = x'$ off Θ_k and $y = y'$ off Θ_k .

We need to bound the expectation $\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [|f(x'_1 \dots x'_j y'_{j+1} \dots y'_n) - f(x'_1 \dots x'_{j-1} y'_j \dots y'_n)|]$ for each site $j \in V$. There are three cases.

- $j \in \Theta_k$. By definition of $\rho_{i,j}^k$ the coupling will yield $x'_j \neq y'_j$ with probability at most $\rho_{i,j}^k$ and so

$$\begin{aligned}
&\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [|f(x'_1 \dots x'_j y'_{j+1} \dots y'_n) - f(x'_1 \dots x'_{j-1} y'_j \dots y'_n)|] \\
&\leq \rho_{i,j}^k \max_{(\sigma,\tau) \in S_j} \{|f(\sigma) - f(\tau)|\} = \rho_{i,j}^k \delta_j(f).
\end{aligned}$$

- $j \notin \Theta_k$ and $j = i$. Since $j \notin \Theta_k$ we have $x_j = x'_j$ and $y_j = y'_j$ so

$$\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [|f(x'_1 \dots x'_j y'_{j+1} \dots y'_n) - f(x'_1 \dots x'_{j-1} y'_j \dots y'_n)|] \leq \delta_j(f) = \delta_i(f).$$

- $j \notin \Theta_k$ and $i \neq j$. In this case we have $x_j = x'_j$ and $y_j = y'_j$ which implies $x'_j = y'_j$ so

$$\mathbf{E}_{(x',y') \in \Psi_k(x,y)} [|f(x'_1 \dots x'_j y'_{j+1} \dots y'_n) - f(x'_1 \dots x'_{j-1} y'_j \dots y'_n)|] = 0.$$

Adding it up we get the statement of the lemma. $\square$

We will use Lemma 6 in conjunction with an inductive proof similar to (V.1.16) in Simon [14] in order to establish the following lemma. It is important to note at this point that the result in Simon is presented for single site heat-bath updates, whereas the following lemma applies to any block dynamics (satisfying the stated assumptions) and weighted sites. This lemma is also a block generalisation of Lemma 11 in Dyer et al. [5].

Lemma 7. Let $\Gamma(k) = \bigcup_{l=1}^k \Theta_l$ then for any $k \in \{1, \dots, m\}$, if $\alpha < 1$ then

$$\Delta(P^{[1]} \dots P^{[k]} f) \leq \alpha \sum_{i \in \Gamma(k)} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f).$$

Proof. Induction on k . Taking $k = 0$ as the base case, we get the definition of Δ .

Assume the statement holds for $k - 1$.

$$\begin{aligned} \Delta(P^{[1]} \dots P^{[k]} f) &\leq \alpha \sum_{i \in \Gamma(k-1)} w_i \delta_i(P^{[k]} f) + \sum_{i \in V \setminus \Gamma(k-1)} w_i \delta_i(P^{[k]} f) \\ &\leq \alpha \sum_{i \in \Gamma(k-1)} \mathbf{1}_{i \notin \Theta_k} w_i \delta_i(f) + \alpha \sum_{i \in \Gamma(k-1)} \sum_{j \in \Theta_k} w_i \rho_{i,j}^k \delta_j(f) \\ &\quad + \sum_{i \in V \setminus \Gamma(k-1)} \mathbf{1}_{i \notin \Theta_k} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k-1)} \sum_{j \in \Theta_k} w_i \rho_{i,j}^k \delta_j(f) \end{aligned}$$

by Lemma 6.

Simplifying and using $\alpha < 1$

$$\begin{aligned} \Delta(P^{[1]} \dots P^{[k]} f) &\leq \alpha \sum_{i \in \Gamma(k-1) \setminus \Theta_k} w_i \delta_i(f) + \sum_{i \in \Gamma(k-1)} \sum_{j \in \Theta_k} w_i \rho_{i,j}^k \delta_j(f) \\ &\quad + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k-1)} \sum_{j \in \Theta_k} w_i \rho_{i,j}^k \delta_j(f) \\ &= \alpha \sum_{i \in \Gamma(k-1) \setminus \Theta_k} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f) \\ &\quad + \sum_{j \in \Theta_k} \delta_j(f) \left(\sum_{i \in \Gamma(k-1)} w_i \rho_{i,j}^k + \sum_{i \in V \setminus \Gamma(k-1)} w_i \rho_{i,j}^k \right) \\ &= \alpha \sum_{i \in \Gamma(k-1) \setminus \Theta_k} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f) + \sum_{j \in \Theta_k} \delta_j(f) \sum_{i \in V} w_i \rho_{i,j}^k \\ &\leq \alpha \sum_{i \in \Gamma(k-1) \setminus \Theta_k} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f) + \sum_{j \in \Theta_k} \delta_j(f) \max_l \sum_{i \in V} w_i \rho_{i,j}^l \\ &\leq \alpha \sum_{i \in \Gamma(k-1) \setminus \Theta_k} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f) + \alpha \sum_{j \in \Theta_k} w_j \delta_j(f) \\ &= \alpha \sum_{i \in \Gamma(k)} w_i \delta_i(f) + \sum_{i \in V \setminus \Gamma(k)} w_i \delta_i(f) \end{aligned}$$

by definition of α . □

Lemma 5 is now a simple consequence of Lemma 7 since

$$\Delta(P_{\rightarrow} f) = \Delta(P^{[1]} \dots P^{[m]} f) \leq \alpha \sum_{i \in V} w_i \delta_i(f) = \alpha \Delta(f)$$

and Theorem 2 follows as discussed above.

3 Application: Edge Scan on an Arbitrary Graph

In this section we prove Theorem 3. That is, we present a general version of a systematic scan on edges and use Theorem 2 to prove that it mixes in $O(\log n)$ scans when $q \geq 2\Delta$. We use $w_i = 1$ for all $i \in V$ and so omit all weights throughout this section. Recall that $\mathcal{M}_{\text{edge}}$ is the systematic scan Markov chain with transition matrix $\prod_{k=1}^m P^{[k]}$ where $\Theta = \{\Theta_k\}_{k=1,\dots,m}$ is an ordered set of edges in G that covers V and $P^{[k]}$ is the transition matrix for performing a heat-bath move on the endpoints of the edge Θ_k .

We need to construct a coupling $\Psi_k(x, y)$ of the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$ for each pair of configurations $(x, y) \in S_i$ that differ only at the colour assigned to site i . Assume without loss of generality that $x_i = 1$ and $y_i = 2$ and also let j and j' be the endpoints of the edge Θ_k . Recall that, since the dynamics uses heat-bath updates, $P^{[k]}(x)$ is the uniform distribution on configurations that agree with x off Θ_k and where no edge containing j or j' is monochromatic. For ease of notation we let $D_1 = P^{[k]}(x)$ and $D_2 = P^{[k]}(y)$. We go on to make the following definitions for $l \in \{1, 2\}$ and $s \in \Theta_k$. $D_l(s)$ is the distribution of the colour assigned to site s induced by D_l , and $[D_l \mid s = c]$ is the uniform distribution on the set of colourings of the sites in Θ_k where site s is assigned colour c . We also let d_l denote the number of configurations with positive measure in D_l and $d_{l,s=c}$ be the number of configurations that assign colour c to site s and have positive measure in D_l .

Definition 8. We will say that the choice $c_1 c_2$ is “valid” for D_l if there is a configuration with positive measure in D_l in which site j is coloured c_1 and site j' is coloured c_2 . Similarly a colour c is “valid” on a site s in D_l if there exists a valid choice for D_l where site s is coloured c .

3.1 Overview of the Coupling

We begin the construction of the coupling $\Psi_k(x, y)$ by giving an overview of the cases we will need to consider and show that they are mutually exclusive and exhaustive of all configurations. It is important to note that, by definition of ρ , the coupling we define may depend on the initial configurations x and y in the sense that if two pairs of configurations (x_1, y_1) and (x_2, y_2) can be distinguished then the couplings $\Psi_k(x_1, y_1)$ and $\Psi_k(x_2, y_2)$ may be defined differently.

First, if i is not adjacent to any site in Θ_k , that is $i \notin \partial\Theta_k$, then $\Psi_k(x, y)$ is the identity coupling where the same colouring is assigned to each distribution. Hence, for $i \notin \partial\Theta_k$ and $j \in \Theta_k$ we have

$$\rho_{i,j}^k = 0.$$

Now suppose that i is adjacent to at least one site in Θ_k , that is $i \in \partial\Theta_k$. We consider the following five cases, which by construction are exhaustive of all possible configurations and mutually exclusive. In the diagrams that relate to these cases a dotted line between a site $j \in \Theta_k$ and a colour 1, say, denotes that no site adjacent to j on the boundary of Θ_k (other than possibly i) is coloured 1. A full line denotes that some site adjacent to j on the boundary of Θ_k (other than possibly i) is coloured 1. The full details of each case of the coupling will be given in section 3.2 along with bounds on $\rho_{i,j}^k$ and $\rho_{i,j'}^k$ where j and j' are the sites included in Θ_k .

1. Exactly one site in Θ_k is adjacent to i . Let this site be labeled j and let the other site in Θ_k be labeled j' . This is shown in Figure 1.
2. Both sites in Θ_k are adjacent to i and no other sites in $\partial\Theta_k$ are coloured 1 or 2. The labeling of the sites in Θ_k is arbitrary. This is shown in Figure 2.

Figure 1: Case 1. Exactly one site in Θ_k is adjacent to i . Let this site be labeled j and let the other site in Θ_k be labeled j' .




Figure 2: Case 2. Both sites in Θ_k are adjacent to i and no other sites in $\partial\Theta_k$ are coloured 1 or 2. The labeling of the sites in Θ_k is arbitrary.




Figure 3: Case 3. Both sites in Θ_k are adjacent to i . One of the sites in Θ_k is adjacent to at least one site, other than i , coloured 1. Let this site be labeled j' . The other site in Θ_k is labeled j and it is not adjacent to any site, other than i , coloured 1 or 2.




Figure 4: Case 4. Both sites in Θ_k are adjacent to i . One of the sites in Θ_k is adjacent to at least one site, other than i , coloured 1 and no sites that are coloured 2. Let this site be labeled j' . The other site in Θ_k , labeled j , is adjacent to at least one site other than i coloured 2 and no sites coloured 1.




Figure 5: Case 5. Both sites in Θ_k are adjacent to i and at least one site, other than i coloured 1. The labeling of the sites in Θ_k is arbitrary.





Figure 6: Case 1. Exactly one site in Θ_k is adjacent to i . Let this site be labeled j and let the other site in Θ_k be labeled j' .



3. Both sites in Θ_k are adjacent to i . One of the sites in Θ_k is adjacent to at least one site, other than i , coloured 1. Let this site be labeled j' . The other site in Θ_k is labeled j and it is not adjacent to any site, other than i , coloured 1 or 2. This is shown in Figure 3.
4. Both sites in Θ_k are adjacent to i . One of the sites in Θ_k is adjacent to at least one site, other than i , coloured 1 and no sites that are coloured 2. Let this site be labeled j' . The other site in Θ_k , labeled j , is adjacent to at least one site other than i coloured 2 and no sites coloured 1. This is shown in Figure 4.
5. Both sites in Θ_k are adjacent to i and at least one site, other than i coloured 1. The labeling of the sites in Θ_k is arbitrary. This is shown in Figure 5.

3.2 Details of Coupling and Proof of Mixing

We will now give the full details of each case of the coupling and establish the required bounds on the influence of site i on sites j and j' . The following lemma is required to establish the coupling for all the stated cases.

Lemma 9. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$. Then for each pair of colours $c_1, c_2 \in C \setminus \{1, 2\}$ the choice $c_1 c_2$ is valid for D_1 if and only if $c_1 c_2$ is valid for D_2 .*

Proof. We start with the *if* direction. Suppose $c_1 c_2$ is valid in D_2 then no site adjacent to j has colour c_1 in D_2 and since $c_1 \neq 1$ no site adjacent to j has colour c_1 in D_1 . Also no site adjacent to j' has colour c_2 in D_2 hence no site adjacent to j' has colour c_2 in D_1 since $c_2 \neq 1$. Since $c_1 c_2$ is valid in D_2 $c_1 \neq c_2$ and so $c_1 c_2$ is valid in D_1 .

The *only if* direction is similar. Suppose $c_1 c_2$ is valid in D_1 then no site adjacent to j has colour c_1 in D_1 and since $c_1 \neq 2$ no site adjacent to j has colour c_1 in D_2 . Also no site adjacent to j' has colour c_2 in D_1 hence no site adjacent to j' has colour c_2 in D_2 again since $c_2 \neq 2$. Since $c_1 c_2$ is valid in D_1 $c_1 \neq c_2$ and so $c_1 c_2$ is valid in D_2 . $\square$

Details of case 1. (Repeated in Figure 6) We construct a coupling $\Psi_k(x, y)$ of the distributions D_1 and D_2 using the following two step process. Let ψ_j be a coupling of $D_1(j)$ and $D_2(j)$ which greedily maximises the probability of assigning the same colour to site j in each distribution. Then, for each pair of colours (c, c') drawn from ψ_j , $\Psi_k(x, y)$ is a coupling, minimising Hamming distance, of the conditional distributions $D_1 \mid j = c$ and $D_2 \mid j = c'$.

Lemma 10. *Let j and j' be the endpoints of an edge Θ_k . If $\{i, j\} \in E(G)$ and $\{i, j'\} \notin E(G)$ then*

$$\rho_{i,j}^k \leq \frac{1}{q - \Delta} \text{ and } \rho_{i,j'}^k \leq \frac{1}{(q - \Delta)^2}.$$

Proof. Assume without loss of generality that $d_1 \geq d_2$, i.e that there are at least as many valid choices for D_1 as for D_2 . Since the only site in Θ_k that is adjacent to site i is j , Lemma 13 of Goldberg, Martin and Paterson [10] lets us upper bound the probability of a discrepancy at site j in a pair of configurations drawn from the coupling $\Psi_k(x, y)$ by assuming that j' is assigned the worst case colour. Now, 1 is not valid for j in D_1 so Lemma 9 implies that only the choice 2 for j in D_1 would cause site j to be assigned a different colour in each configuration drawn from the coupling. Now observe that site j has at most $\Delta - 1$ neighbours (excluding j') and each of them could invalidate one colour choice for j in both distributions. If j' is assigned a colour not already adjacent to j then j is adjacent to at most Δ sites each assigned a different colour, leaving at least $q - \Delta$ valid colours for j in D_1 and so the probability of assigning 2 to j in D_1 during step 1 of the coupling is at most $\frac{1}{q - \Delta}$ since the coupling is greedy. This establishes the bound on $\rho_{i,j}^k$ since

$$\rho_{i,j}^k = \max_{(x,y) \in S_i} \{\Pr_{(x',y') \in \Psi_k(x,y)}(x'_j \neq y'_j)\} \leq \frac{1}{q - \Delta}.$$

Now from the definition of the coupling it follows easily that if the same colour, c , is assigned to site j in each distribution during the first step of the coupling then the colour assigned to site j' in the second step will be the same in each distribution since the conditional distributions $D_1 \mid j = c$ and $D_2 \mid j = c$ are the same. If different colours are assigned to j in each distribution then the second step of the coupling is simply the case of colouring a single site adjacent to exactly one discrepancy. The argument from above says that at most one colour assigned to j' in D_1 will cause a discrepancy at site j' in the coupling and also that there are at least $q - \Delta$ valid choices for j' in D_1 . Hence we have $\max_{(x,y) \in S_i} \{\Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'} \mid x'_j = c, y'_j = c')\} \leq \frac{1}{q - \Delta}$ and so

$$\begin{aligned} \rho_{i,j'}^k &= \max_{(x,y) \in S_i} \{\Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'})\} \\ &= \max_{(x,y) \in S_i} \left\{ \sum_{c,c'} \Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'} \mid x'_j = c, y'_j = c') \Pr_{(x',y') \in \Psi_k(x,y)}(x'_j = c, y'_j = c') \right\} \\ &\leq \frac{1}{q - \Delta} \max_{(x,y) \in S_i} \left\{ \sum_{c,c'} \Pr_{(x',y') \in \Psi_k(x,y)}(x'_j = c, y'_j = c') \right\} \\ &\leq \frac{1}{(q - \Delta)^2} \end{aligned}$$

using the bound from $\rho_{i,j}^k$ which completes the proof. $\square$

The following lemmas are required to define the coupling and bound the influence of a site $i \in \partial\Theta_k$ on sites j and j' when i is adjacent to both sites j and j' .

Lemma 11. *Let j and j' be the endpoints of an edge and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$. If 1 is valid for j in D_2 and 2 is valid for j in D_1 then the choice $2c_2$ is valid in D_1 if and only if $1c_2$ is valid in D_2 .*

Proof. Suppose that $2c_2$ is valid in D_1 then $c_2 \in C \setminus \{1, 2\}$ since i is adjacent to j' (and $x_i = 1$). Since 1 is valid for j in D_2 it follows that $1c_2$ is valid in D_2 since the only colour adjacent to j' in D_2 that is (possibly) not adjacent to j' in D_1 is 2, but $c_2 \neq 2$.

For the reverse direction suppose that $1c_2$ is valid in D_2 . Then $c_2 \in C \setminus \{1, 2\}$ since i is adjacent to j' . Since 2 is valid for j in D_1 it follows that $2c_2$ is valid in D_1 since the only colour adjacent to j' in D_1 that is (possibly) not adjacent to j' in D_2 is 1, but $c_2 \neq 1$. $\square$

Lemma 12. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$. If 1 is valid for j' in D_2 and 2 is valid for j' in D_1 then the choice $c_1 2$ is valid in D_1 if and only if $c_1 1$ is valid in D_2 .*

Proof. Suppose that $c_1 2$ is valid in D_1 then $c_1 \in C \setminus \{1, 2\}$ since i is adjacent to j' . Since 1 is valid for j' in D_2 $c_1 1$ is valid in D_2 since the only colour adjacent to j in D_2 that is (possibly) not adjacent to j in D_1 is 2, but $c_1 \neq 2$.

Also, suppose that $c_1 1$ is valid in D_2 then $c_1 \in C \setminus \{1, 2\}$ since i is adjacent to j' . Since 2 is valid for j' in D_1 $c_1 2$ is valid in D_1 since the only colour adjacent to j in D_1 that is (possibly) not adjacent to j in D_2 is 1, but $c_1 \neq 1$. $\square$

Lemma 13. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$.*

(i) *Suppose that 1 is valid for j in D_2 . For all $c \in C$ where c is valid for j in D_2 , if 1 is valid for j' in D_2 then*

$$d_{2,j=1} \leq d_{2,j=c} \leq d_{2,j=1} + 1$$

else

$$d_{2,j=1} - 1 \leq d_{2,j=c} \leq d_{2,j=1}.$$

(ii) *Suppose that 2 is valid for j in D_1 . For all $c \in C$ where c is valid for j in D_1 , if 2 is valid for j' in D_1 then*

$$d_{1,j=2} \leq d_{1,j=c} \leq d_{1,j=2} + 1$$

else

$$d_{1,j=2} - 1 \leq d_{1,j=c} \leq d_{1,j=2}.$$

Proof. Part (i). Consider some valid colour c other than 1 for j in D_2 . For each valid choice $1c_2$ for D_2 the choice cc_2 is also valid for D_2 except when $c = c_2$. If 1 is valid for j' in D_2 then the choice $c1$ is also valid for D_2 .

Now consider some invalid choice $1c_2$ for D_2 where $c_2 \neq 1$. Since $1c_2$ is not valid for D_2 it follows that c_2 is not valid for j' in D_2 and hence no more choices can be valid for D_2 , which guarantees the upper bounds.


Part (ii) is similar. Consider some valid colour c other than 2 for j in D_1 . For each valid choice $2c_2$ for D_1 the choice cc_2 is also valid for D_1 except when $c = c_2$. If 2 is valid for j' in D_1 then the choice $c2$ is also valid for D_1 .

Finally consider some invalid choice $2c_2$ for D_1 where $c_2 \neq 2$. Since $2c_2$ is not valid for D_1 it follows that c_2 is not valid for j' in D_1 and hence no more choices can be valid for D_1 , which guarantees the upper bounds. $\square$

We are now ready to define the coupling for the remaining cases.

Details of case 2. (Repeated in Figure 7) We construct the $\Psi_k(x, y)$ of the distributions D_1

Figure 7: Case 2. Both sites in Θ_k are adjacent to i and no other sites in $\partial\Theta_k$ are coloured 1 or 2. The labeling of the sites in Θ_k is arbitrary.



and D_2 as follows. For each valid choice of the form c_1c_2 for D_1 where $c_1 \neq 2$ and $c_2 \neq 2$ Lemma 9 guarantees that c_1c_2 is valid for D_2 so we let

$$\Pr_{(x',y') \in \Psi_k(x,y)}(x' = y' = c_1c_2) = \frac{1}{d_1}.$$

For each valid choice of the form $2c_2$ in D_1 the choice $1c_2$ is valid in D_2 by Lemma 11 so we let

$$\Pr_{(x',y') \in \Psi_k(x,y)}(x' = 2c_2, y' = 1c_2) = \frac{1}{d_1}. \quad (3)$$

Lemma 11 also guarantees that there are no remaining valid choices for D_2 of the form $1c_2$. Finally for each valid choice c_12 for D_1 the choice c_11 is valid in D_2 by Lemma 12 so let

$$\Pr_{(x',y') \in \Psi_k(x,y)}(x' = c_12, y' = c_11) = \frac{1}{d_1} \quad (4)$$

which completes the coupling since $d_1 = d_2$ and all the probability in both D_1 and D_2 has hence been used.

Lemma 14. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$. If 2 is valid for both j and j' in D_1 and 1 is valid for both j and j' in D_2 then*

$$\rho_{i,j}^k \leq \frac{1}{q - \Delta + 1} \text{ and } \rho_{i,j'}^k \leq \frac{1}{q - \Delta}.$$


Proof. This is case 2 of the coupling. Note from Lemma 11 that $d_{1,j=2} = d_{2,j=1}$ so for ease of reference let $d = d_{1,j=2} = d_{2,j=1}$ and let $d' = d_{1,j'=2} = d_{2,j'=1}$ by Lemma 12. Also let $s = \sum_c d_{2,j=c} - d - d'$ which is the number of valid choices for D_2 other than choices of the form $1c_2$ and c_11 . Note that the number of valid choices for D_1 is $d_1 = s + d + d'$.

As there are no restrictions on colours assigned to the sites in $\partial\Theta_k \setminus \{i\}$ each of the neighbours of j could be assigned a different colour, and the same is true for the neighbours of j' . Hence we get the following lower-bounds on d and d' :

$$q - \Delta \leq d \text{ and } q - \Delta \leq d'.$$

To lower bound s observe that $s = \sum_c d_{2,j=c} - d - d' = \sum_{c \neq 1} d_{2,j=c} - d'$. Let $J \subseteq C \setminus \{1\}$ be the set of colours, excluding 1, that are valid for j in D_2 . By definition of d' , at least d' colours

Figure 8: Case 3. Both sites in Θ_k are adjacent to i . One of the sites in Θ_k is adjacent to at least one site, other than i , coloured 1. Let this site be labeled j' . The other site in Θ_k is labeled j and it is not adjacent to any site, other than i , coloured 1 or 2.



other than 1 must be valid for site j in D_2 so the size of J is at least d' . Since 1 is valid for j' in D_2 we use the lower bound on $d_{2,j=c}$ from Lemma 13 (i) and hence

$$\begin{aligned} s &= \sum_{c \in J} d_{2,j=c} - d' \\ &\geq d' \min_{c \in J} \{d_{2,j=c}\} - d' \\ &\geq d'd - d'. \end{aligned}$$

From the coupling, j will be assigned a different colour in each distribution whenever a choice of the form $2c_2$ is made for D_1 . From (3) this happens with probability $\frac{d}{d_1} = \frac{d}{d+d'+s}$ since d is the number of valid choices for D_1 of the form $2c_2$. Similarly from (4), j' will become a discrepancy in the coupling whenever a choice of the form c_12 is made for D_1 , which happens with probability $\frac{d'}{d+d'+s}$. Hence

$$\rho_{i,j}^k \leq \frac{d}{d+d'+s} \text{ and } \rho_{i,j'}^k \leq \frac{d'}{d+d'+s}.$$

Starting with $\rho_{i,j}^k$

$$\rho_{i,j}^k \leq \frac{d}{d+d'+s} \leq \frac{d}{d+dd'} \leq \frac{1}{d'+1} \leq \frac{1}{q-\Delta+1}$$

using the lower bounds of s and d' . Similarly using the lower bounds of s and d

$$\rho_{i,j'}^k \leq \frac{d'}{d+d'+s} \leq \frac{d'}{d+dd'} \leq \frac{1}{d} \leq \frac{1}{q-\Delta}$$


which implies the statement of the lemma. $\square$

Details of case 3. (Repeated in Figure 8) We construct the coupling $\Psi_k(x, y)$ of D_1 and D_2 using the following two step process. Let Ψ_j be a coupling of $D_1(j')$ and $D_2(j')$ which greedily maximises the probability of assigning the same colour to site j' in each distribution. Then for each pair of colours (c, c') drawn from Ψ_j we complete $\Psi_k(x, y)$ by letting it be the coupling, greedily minimising Hamming distance, of the conditional distributions $D_1 \mid j' = c$ and $D_2 \mid j' = c'$.

Lemma 15. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$. If 2 is valid for j in D_1 , 1 is valid for j in D_2 and 1 is not valid for j' in D_2 then*

$$\rho_{i,j'}^k \leq \frac{1}{q-\Delta+1} \text{ and } \rho_{i,j}^k \leq \frac{1}{q-\Delta}.$$

Figure 9: The pair of configurations after the colour of site j' has been assigned during the first step of the coupling.



Proof. This is case 3 of the coupling. Note from Lemma 11 that $d_{1,j=2} = d_{2,j=1}$ and let $s = \sum_c d_{2,j=c} - d_{2,j=1} = \sum_{c \neq 1} d_{2,j=c}$ denote the number of valid choices for D_2 other than choices of the form $1c_2$. The number of valid choices for D_1 is then $d_1 = s + d_{1,j=2} + d_{1,j'=2}$.

Since 1 is not valid for j' in D_2 at least one site other than i on the boundary of Θ_k must be coloured 1 in D_1 (we say that some site s on the boundary of Θ_k is coloured c in D_1 if there exists a configuration with positive measure in D_1 in which site s is coloured c). As there are no restrictions on the neighbourhood of j each neighbour of j may be assigned a different colour in D_1 . Hence we get the following lower bounds on $d_{1,j=2}$ and $d_{1,j'=2}$

$$q - \Delta + 1 \leq d_{1,j=2} \text{ and } q - \Delta \leq d_{1,j'=2}.$$

To lower bound s observe that exactly $d_{1,j'=2}$ colours other than 1 are valid for site j in D_2 and let J be the set of colours, excluding 1, that are valid for j in D_2 , then

$$s = \sum_{c \in J} d_{2,j=c} \geq d_{1,j'=2} \min_{j \in J} \{d_{2,j=c}\} \geq d_{1,j'=2} (d_{1,j=2} - 1)$$

where we used the bound $d_{1,j=2} - 1 \leq d_{1,j=2}$ for $c \in J$ from Lemma 13 (i) since 1 is not valid for j' in D_2 .

We consider $\rho_{i,j'}^k$ first. Suppose that a choice of the form c_1c_2 is valid for D_2 , in which case $c_1 \neq 2$ and $c_2 \notin \{1, 2\}$ by the conditions of case 3 of the coupling. Firstly if $c_1 \neq 1$ then c_1c_2 is also valid for D_1 by Lemma 9. If $c_1 = 1$ then the choice $2c_2$ is valid for D_1 by Lemma 11 and hence $d_1 \geq d_2$. Note in particular that if a choice c_1c_2 where $c_2 \neq 2$ is valid for D_1 then it is also valid for D_2 . Therefore, a different colour will only be assigned to site j' in each distribution if j' is coloured 2 in D_1 during the first step of the coupling since the Hamming distance at site j' is minimised greedily. There are $d_{1,j'=2}$ colourings assigning 2 to j' in D_1 and hence

$$\rho_{i,j'}^k \leq \frac{d_{1,j'=2}}{d_{1,j=2} + d_{1,j'=2} + s} \leq \frac{d_{1,j'=2}}{d_{1,j=2} + (1 + d_{1,j'=2})} \leq \frac{1}{d_{1,j=2}} \leq \frac{1}{q - \Delta + 1}$$

using the lower bounds on s and $d_{1,j=2}$.

Now consider $\rho_{i,j}^k$. Suppose that (c'_1, c'_2) is the pair of colours drawn for site j' in the first step of the coupling. The second step of $\Psi_k(x, y)$ then couples the conditional distributions $D_1 \mid j' = c'_1$ and $D_2 \mid j' = c'_2$ greedily to minimise Hamming distance. First suppose that $c'_1 \neq c'_2$. It was pointed out in the analysis above that if $c'_1 \neq c'_2$ then $c'_1 = 2$ and the resulting configuration is shown in Figure 9. We make the following observations about the resulting conditional distributions $D_1 \mid j' = 2$ and $D_2 \mid j' = c'_2$.

- The colour 2 is *not* valid for j in either distribution $D_1 \mid j' = 2$ or $D_2 \mid j' = c'_2$.

- The colour 1 is *not* valid for j in $D_1 \mid j' = 2$ but *could* be valid for j in $D_2 \mid j' = c'_2$.
- The colour c'_2 *could* be valid for j in $D_1 \mid j' = 2$ but is *not* valid for j in $D_2 \mid j' = c'_2$.
- For each $c \in C \setminus \{1, 2, c'_2\}$ the colour c is valid for j in $D_1 \mid j' = 2$ *if and only if* c is valid for j in $D_2 \mid j' = c'_2$.

These observations show that this case is a single-site disagreement sub problem and that there must be at least $(q - 3) - (\Delta - 2) = q - \Delta - 1$ colours that are valid for j in both conditional distributions since j has at most $\Delta - 2$ neighbours other than i and j' . Also, there is at most one colour which is valid for j in one distribution but not in the other and since the coupling greedily maximises Hamming distance this implies

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j \mid x'_{j'} \neq y'_{j'}) \leq \frac{1}{q - \Delta}.$$

Now suppose that the same colour c , say, is drawn for site j' in both distributions during the first step of the coupling. Then the only site adjacent to i that is coloured differently in the conditional distributions $D_1 \mid j' = c$ and $D_2 \mid j' = c$ is site i , so using a similar reasoning to above we find

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j \mid x'_{j'} = y'_{j'}) \leq \frac{1}{q - \Delta}$$

and thus

$$\begin{aligned} \rho_{i,j}^k &= \max_{(x, y) \in S_i} \{ \Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j) \} \\ &= \max_{(x, y) \in S_i} \{ \Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j \mid x'_{j'} \neq y'_{j'}) \Pr_{(x', y') \in \Psi_k(x, y)}(x'_{j'} \neq y'_{j'}) \\ &\quad + \Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j \mid x'_{j'} = y'_{j'}) \Pr_{(x', y') \in \Psi_k(x, y)}(x'_{j'} = y'_{j'}) \} \\ &\leq \max_{(x, y) \in S_i} \left\{ \frac{1}{q - \Delta} \Pr_{(x', y') \in \Psi_k(x, y)}(x'_{j'} \neq y'_{j'}) + \frac{1}{q - \Delta} \Pr_{(x', y') \in \Psi_k(x, y)}(x'_{j'} = y'_{j'}) \right\} \\ &= \frac{1}{q - \Delta} \max_{(x, y) \in S_i} \{ \Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j) + \Pr_{(x', y') \in \Psi_k(x, y)}(x'_j = y'_j) \} = \frac{1}{q - \Delta} \end{aligned}$$

which completes the proof. $\square$


Details of case 4. (Repeated in Figure 10) We assume without loss of generality that $d_1 \geq d_2$ and construct the coupling $\Psi_k(x, y)$ of D_1 and D_2 as follows. For each valid choice of the form $c_1 c_2$ for D_1 where $c_1 \neq 1$ and $c_2 \neq 2$ Lemma 9 guarantees that $c_1 c_2$ is also valid for D_2 so we construct $\Psi_k(x, y)$ such that

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x' = y' = c_1 c_2) = \frac{1}{d_1}.$$

This leaves the set $Z_1 = \{c_1 2 \mid c_1 2 \text{ valid in } D_1\}$ of valid choices for D_1 and $Z_2 = \{1 c_2 \mid 1 c_2 \text{ valid in } D_2\} \subseteq D_2$ for D_2 . Observe that $z_1 \geq z_2$ where z_1 and z_2 denote the size of Z_1 and Z_2 respectively. Let $Z_1(t)$ denote the t -th element of Z_1 and similarly for Z_2 . Then for $1 \leq t \leq z_2$ let

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x' = Z_1(t), y' = Z_2(t)) = \frac{1}{d_1}$$

Figure 10: Case 4. Both sites in Θ_k are adjacent to i . One of the sites in Θ_k is adjacent to at least one site, other than i , coloured 1 and no sites that are coloured 2. Let this site be labeled j' . The other site in Θ_k , labeled j , is adjacent to at least one site other than i coloured 2 and no sites coloured 1.



and for each pair $z_2 + 1 \leq t \leq z_2$ and $h \in D_2$ let

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x' = Z_1(t), y' = h) = \frac{1}{d_1 d_2}.$$

It is easy to verify that each valid colouring has the correct weight in $\Psi_k(x, y)$ so this completes the coupling.

Lemma 16. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$. If 1 is valid for j in D_2 , 1 is not valid for j' in D_2 and 2 is not valid for j in D_1 then*

$$\rho_{i,j}^k \leq \rho_{i,j'}^k \leq \frac{1}{q - \Delta}.$$

Proof. This is case 4 of the coupling. Let $s = \sum_c d_{2,j=c} - d_{2,j=1}$ be the number of valid choices for D_2 other than choices of the form $1c_2$. Observe that $d_2 = s + d_{1,j'=2}$ and note that $d_{1,j'=2} \geq d_{2,j=1}$ since we have assumed $d_1 \geq d_2$ in the construction of the coupling. At least one neighbour, other than i , of j' on the boundary of Θ_k is coloured 1 in D_1 and we get the following lower-bound on $d_{2,j=1}$ since all other neighbours of j' may be assigned a different colour

$$q - \Delta + 1 \leq d_{2,j=1}.$$

We bound s using the same argument as in the proof of Lemma 15 and get

$$d_{1,j'=2}(d_{2,j=1} - 1) \leq s.$$


Since 2 is not valid for j in D_1 the first $d_{2,j=1}$ choices of the form $c_1 2$ for D_1 are matched with some choice of the form $1c_1$ for D_2 with probability $1/d_1$ resulting in a different colour being assigned to both sites j and j' in each distribution. Each of the $d_{1,j'=2} - d_{2,j=1}$ remaining valid choices for D_1 is matched with each valid choice for D_2 with probability $\frac{1}{d_1 d_2}$ resulting in a disagreement at j' (since 2 is not valid for j' in D_2) and potentially also at j so $\rho_{i,j}^k \leq \rho_{i,j'}^k$. Hence the probability of making a choice of the form $c_1 2$ for D_1

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x'_{j'} = 2) = \frac{d_{2,j=1}}{d_1} + \sum_{h \in D_2} \frac{d_{1,j'=2} - d_{2,j=1}}{d_1 d_2} = \frac{d_{1,j'=2}}{d_1}$$

is an upper bound on the disagreement probabilities at both sites j and j' . Using the lower bounds on s and $d_{2,j=1}$ we have

$$\rho_{i,j}^k \leq \rho_{i,j'}^k \leq \frac{d_{1,j'=2}}{d_1} = \frac{d_{1,j'=2}}{d_{1,j'=2} + s} \leq \frac{d_{1,j'=2}}{d_{1,j'=2} + (d_{2,j=1} - 1)d_{1,j'=2}} \leq \frac{1}{q - \Delta}$$

Figure 11: Case 5. Both sites in Θ_k are adjacent to i and at least one site, other than i coloured 1. The labeling of the sites in Θ_k is arbitrary.



which completes the proof. $\square$

Details of case 5. (Repeated in Figure 11) First observe that 1 is not valid for both j and j' in either distribution D_1 or D_2 so $d_1 = d_2 + d_{1,j=2} + d_{1,j'=2} \geq d_2$ by Lemma 9, since any choice valid for D_2 does not assign colour 2 to any site in Θ_k . Let Z_1 and Z_2 be the sets of colourings valid for D_1 and D_2 respectively. We define the following mutually exclusive subsets of Z_1 . $Z_j = \{2c_2 \mid 2c_2 \in Z_1\}$, $Z_{j'} = \{c_1 2 \mid c_1 2 \in Z_1\}$ and $Z = Z_1 \setminus (Z_j \cup Z_{j'}) = Z_2$. By construction, the union of these three subsets is Z_1 and note that the size of Z_j is $d_{1,j=2}$, the size of $Z_{j'}$ is $d_{1,j'=2}$ and the size of Z is d_2 .

First we consider choices from Z for D_1 . For each choice $h \in Z$ we have $h \in Z_2$ by construction of Z and so we use the identity coupling and let

$$\Pr_{(x',y') \in \Psi_k(x,y)}(x' = y' = h) = \frac{1}{d_1}.$$

We let the remainder of the coupling minimise Hamming distance. First consider the choices for D_1 in Z_j . We construct $\Psi_k(x, y)$ such that it minimises Hamming distance and assigns probability $1/d_1$ to each choice for D_1 in Z_j whilst ensuring that for each choice $g \in Z_2$ for D_2

$$\sum_{h \in Z_j} \Pr_{(x',y') \in \Psi_k(x,y)}(x' = h, y' = g) = \frac{d_{1,j=2}}{d_1 d_2}.$$

Similarly we assign probability $1/d_1$ to each choice for D_1 in $Z_{j'}$ whilst also requiring that for each choice $g \in Z_2$ for D_2

$$\sum_{h \in Z_{j'}} \Pr_{(x',y') \in \Psi_k(x,y)}(x' = h, y' = g) = \frac{d_{1,j'=2}}{d_1 d_2}.$$

To see that this ensures that the coupling is fair observe that each choice $h \in Z_1$ receives weight $1/d_1$ and each choice $g \in Z_2$ weight

$$\frac{1}{d_1} + \frac{d_{1,j=2}}{d_1 d_2} + \frac{d_{1,j'=2}}{d_1 d_2} = \frac{d_2 + d_{1,j=2} + d_{1,j'=2}}{d_1 d_2} = \frac{1}{d_2}$$

since $d_2 + d_{1,j=2} + d_{1,j'=2} = d_1$.

Remark. Note that a coupling satisfying these requirements always exists. We will not give the detailed construction of $\Psi_k(x, y)$ here, but in the subsequent proof we will consider three cases. In the first two cases *any* coupling minimising Hamming distance will be sufficient to establish the required bounds on the influence of i on j . In the final case we will need a detailed construction of the coupling and so will provide it together with the proof for ease of reference.

Lemma 17. *Let j and j' be the endpoints of an edge Θ_k and suppose that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$. If 1 is not valid for j in D_2 and 1 is not valid for j' in D_2 then*

$$\rho_{i,j}^k \leq \frac{1}{q - \Delta + 1} + \frac{1}{(q - \Delta + 1)^2} \text{ and } \rho_{i,j'}^k \leq \frac{1}{q - \Delta + 1} + \frac{1}{(q - \Delta + 1)^2}.$$

Proof. This is case 5 of the coupling. We consider three separate cases. Firstly suppose that 2 is not valid for both j and j' in D_1 . Then the only valid choices for D_1 are of the form $c_1 c_2$ where $c_1, c_2 \in C \setminus \{1, 2\}$ and each such choice is also valid in D_2 as observed in the construction of the coupling. The same colouring is selected for each distribution and hence

$$\rho_{i,j}^k = 0 \text{ and } \rho_{i,j'}^k = 0.$$

Next suppose that exactly one site in Θ_k , j' say, is adjacent to some site coloured 2 in D_1 . As in the previous case, each choice that is valid in both D_1 and D_2 is matched using the identity matching and does not cause a discrepancy at any site. However if a choice of the form $2c$ is made for D_1 then site j will be coloured differently in each colouring drawn from $\Psi_k(x, y)$ and the colour at site j' may also be different so $\rho_{i,j'}^k \leq \rho_{i,j}^k$. Since all choices of the form $c2$ are not valid for D_1 , making a choice of the form $2c$ for D_1 is the only way to create a disagreement at any site in the coupling and so

$$\rho_{i,j'}^k \leq \rho_{i,j}^k \leq \frac{d_{1,j=2}}{d_1}$$

since $d_{1,j=2}$ is the number of valid choices for D_1 of the form $2c$. We need to establish a lower bound of d_1 and observe that, for c valid for j in D_1 , $d_{1,j=2} - 1 \leq d_{1,j=c}$ by Lemma 13 (ii) since 2 is not valid for j' in D_1 . Let v be the number of colours that are valid for site j in D_1 . Then v is lower bounded by $q - \Delta + 2 \leq v$ since at least two of the sites (including i) adjacent to j on the boundary of Θ_k are coloured 1 in D_1 . Also, since at least one site (other than j and i) adjacent to j' is coloured 1 and another is coloured 2 in D_1 , we have $q - \Delta + 2 \leq d_{1,j=2}$. Using the lower bounds on v and $d_{1,j=c}$ we have, letting J denote the set of colours other than 2 that are valid for j in D_1 ,

$$\begin{aligned} d_1 &= \sum_c d_{1,j=c} = d_{1,j=2} + \sum_{c \in J} d_{1,j=c} \geq d_{1,j=2} + \sum_{c \in J} (d_{1,j=2} - 1) \\ &\geq (v - 1)(d_{1,j=2} - 1) + d_{1,j=2} \geq (q - \Delta + 2)d_{1,j=2} - (q - \Delta + 1) \end{aligned}$$

and hence using the lower bound on $d_{1,j=2}$

$$\frac{1}{\rho_{i,j}^k} \geq \frac{(q - \Delta + 2)d_{1,j=2} - (q - \Delta + 1)}{d_{1,j=2}} \geq q - \Delta + 2 - \frac{q - \Delta + 1}{q - \Delta + 2} > q - \Delta + 1$$

which gives the bounds required by the statement of the lemma.

Finally consider the case when the colour 2 is valid for both j and j' in D_1 . In this case we will provide details of the construction of $\Psi_k(x, y)$ when required. We begin by establishing some required bounds. Since 1 is not valid for j' in D_2 at least two neighbours of j' (including i) must be coloured 1 in D_1 and the same applies to the neighbourhood of j , so we get the following lower bounds on $d_{1,j=2}$ and $d_{1,j'=2}$

$$q - \Delta + 1 \leq d_{1,j=2} \text{ and } q - \Delta + 1 \leq d_{1,j'=2}.$$

We also require bounds on $d_{2,j=c}$ and $d_{2,j'=c}$ for other colours c . Suppose that the choice cc' is valid in D_2 then, since $c, c' \in C \setminus \{1, 2\}$, cc' is also valid for D_1 by Lemma 9. Furthermore, the choice $c2$ is valid in D_1 (but not D_2) so $d_{1,j=c} - 1 = d_{2,j=c}$. Lemma 13 (ii) guarantees that $d_{1,j=2} \leq d_{1,j=c} \leq d_{1,j=2} + 1$ so

$$d_{1,j=2} - 1 \leq d_{2,j=c} \leq d_{1,j=2}$$

for any c valid for j in D_1 and a similar argument gives the bound

$$d_{1,j'=2} - 1 \leq d_{2,j'=c} \leq d_{1,j'=2}$$

for any colour c valid for j' in D_2 . Observe that exactly $d_{1,j'=2}$ colours must be valid for site j in D_2 so using the stated bounds on $d_{2,j=c}$ we have the following bounds on d_2

$$d_{1,j'=2}(d_{1,j=2} - 1) \leq d_2 \leq d_{1,j'=2}d_{1,j=2}.$$

We bound the probability of disagreements at sites j and j' from choices made for D_1 . From the coupling we again note that if a choice c_1c_2 where $c_1 \neq 2$ and $c_2 \neq 2$ is made for D_1 then there will be no disagreements at any site in Θ_k .

Consider making a valid choice of the form $2c$ for D_1 . Firstly, such a choice for D_1 will cause site j to be coloured differently in any pair of colourings drawn from the coupling since 2 is not valid for j in D_2 . We construct $\Psi_k(x, y)$ such that the choice $2c$ for D_1 is matched with a choice of the form $c'c$ for D_2 as long as such a choice that has not exceeded its aggregated probability exists. Let J denote the set of choices of the form $c'c$ that are valid for D_2 and note that the size of J is $d_{2,j'=c}$. The total aggregated weight of all choices of the form $c'c$ for D_2 is

$$\sum_{g \in J} \sum_{h \in Z_1} \Pr_{(x', y') \in \Psi_k(x, y)}(x' = h, y' = g) = \sum_{g \in J} \frac{d_{1,j=2}}{d_1 d_2} = \frac{d_{2,j'=c} d_{1,j=2}}{d_1 d_2}$$

so as long as

$$\frac{1}{d_1} \leq \frac{d_{2,j'=c} d_{1,j=2}}{d_1 d_2}$$

there is enough probability available in Z_2 to match all the weight of the choice $2c$ for D_1 with a choice of the form $c'c$ for D_2 and hence assigning the same colour, c , to site j' in any pair of colourings drawn from the coupling. If there is not enough unassigned weight available in Z_2 then the coupling will match at much probability as possible, $\frac{d_{2,j'=c} d_{1,j=2}}{d_1 d_2}$, with choices of the form $c'c$ for Z_2 but the remaining probability will be matched with choices not assigning colour c to site j' in Z_2 . Hence we obtain the following probabilities conditioned on making a choice of the form $2c$ for D_1 .

$$\Pr_{(x', y') \in \Psi_k(x, y)}(x'_j \neq y'_j \mid x' = 2c) = 1$$

and

$$\begin{aligned} \Pr_{(x', y') \in \Psi_k(x, y)}(x'_{j'} \neq y'_{j'} \mid x' = 2c) &\leq \max \left(0, 1 - \frac{d_{2,j'=c} d_{1,j=2}}{d_2} \right) \\ &\leq \max \left(0, 1 - \frac{(d_{1,j'=2} - 1) d_{1,j=2}}{d_{1,j=2} d_{1,j'=2}} \right) \\ &\leq \frac{1}{d_{1,j'=2}} \end{aligned}$$

using the bounds on d_2 and $d_{1,j'=c}$. Lastly observe that there are $d_{1,j=2}$ valid choices for D_1 of the form $2c$ so

$$\sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x' = 2c) = \frac{d_{1,j=2}}{d_1} = \frac{d_{1,j=2}}{d_{1,j=2} + d_{1,j'=2} + d_2}.$$

Also consider making a valid choice of the form $c2$ for D_1 . This case is symmetric to the construction above, but we include it for completeness. A choice of the form $c2$ for D_1 will cause site j' to be coloured differently in any pair of colourings drawn from the coupling since 2 is not valid for j' in D_2 . We hence construct $\Psi_k(x,y)$ such that it matches the choice $c2$ for D_1 with a choice of the form cc' for D_2 as long as such a choice that has not exceeded its aggregated probability exists. Let J' denote the set of choices of the form cc' that are valid for D_2 and note that the size of J' is $d_{2,j=c}$. The total aggregated weight of all choices of the form cc' for D_2 is

$$\sum_{g \in J'} \sum_{h \in Z_1} \Pr_{(x',y') \in \Psi_k(x,y)}(x' = h, y' = g) = \sum_{g \in J'} \frac{d_{1,j'=2}}{d_1 d_2} = \frac{d_{2,j=c} d_{1,j'=2}}{d_1 d_2}$$

so as long as

$$\frac{1}{d_1} \leq \frac{d_{2,j=c} d_{1,j'=2}}{d_1 d_2}$$

there is enough weight available in Z_2 to match all the weight of the choice $c2$ for D_1 with a choice of the form cc' for D_2 and hence assigning the same colour, c , to site j in any pair of colourings drawn from the coupling. If there is not enough unassigned weight available in Z_2 then the coupling will match at much weight as possible, $\frac{d_{2,j=c} d_{1,j'=2}}{d_1 d_2}$, with choices of the form cc' for Z_2 but the remaining weight will be matched with choices not assigning colour c to site j in Z_2 . Hence we obtain the following probabilities conditioned on making a choice of the form $c2$ for D_1

$$\begin{aligned} \Pr_{(x',y') \in \Psi_k(x,y)}(x'_j \neq y'_j \mid x' = c2) &\leq \max\left(0, 1 - \frac{d_{2,j=c} d_{1,j'=2}}{d_2}\right) \\ &\leq \max\left(0, 1 - \frac{d_{1,j'=2}(d_{1,j=2} - 1)}{d_{1,j=2} d_{1,j'=2}}\right) \\ &\leq \frac{1}{d_{1,j=2}} \end{aligned}$$

using the bounds on d_2 and $d_{1,j=c}$, and as before we also have

$$\Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'} \mid x' = c2) = 1 \text{ and } \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) = \frac{d_{1,j'=2}}{d_{1,j=2} + d_{1,j'=2} + d_2}.$$

Using the conditional probabilities and the bounds on d_2 , $d_{1,j=2}$ and $d_{1,j'=2}$ we find

$$\begin{aligned}
\rho_{i,j}^k &= \max_{(x,y) \in S_i} \{ \Pr_{(x',y') \in \Psi_k(x,y)}(x'_j \neq y'_j) \} \\
&= \max_{(x,y) \in S_i} \left\{ \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x'_j \neq y'_j \mid x' = 2c) \Pr_{(x',y') \in \Psi_k(x,y)}(x' = 2c) \right. \\
&\quad \left. + \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x'_j \neq y'_j \mid x' = c2) \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \sum_c \left[\Pr_{(x',y') \in \Psi_k(x,y)}(x' = 2c) + \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) \frac{1}{d_{1,j=2}} \right] \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \frac{d_{1,j=2}}{d_{1,j=2} + d_{1,j'=2} + d_2} + \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) \frac{1}{d_{1,j=2}} \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \frac{d_{1,j=2}}{d_{1,j=2}(1 + d_{1,j'=2})} + \frac{d_{1,j'=2}}{d_{1,j=2} + d_{1,j'=2} + d_2} \frac{1}{d_{1,j=2}} \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \frac{1}{1 + d_{1,j'=2}} + \frac{d_{1,j'=2}}{d_{1,j=2}(1 + d_{1,j'=2})} \frac{1}{d_{1,j=2}} \right\} \\
&\leq \frac{1}{q - \Delta + 2} + \frac{1}{(q - \Delta + 1)^2}
\end{aligned}$$

and again by symmetry

$$\begin{aligned}
\rho_{i,j'}^k &= \max_{(x,y) \in S_i} \{ \Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'}) \} \\
&= \max_{(x,y) \in S_i} \left\{ \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'} \mid x' = 2c) \Pr_{(x',y') \in \Psi_k(x,y)}(x' = 2c) \right. \\
&\quad \left. + \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x'_{j'} \neq y'_{j'} \mid x' = c2) \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \sum_c \left[\Pr_{(x',y') \in \Psi_k(x,y)}(x' = 2c) + \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) \frac{1}{d_{1,j'=2}} \right] \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \frac{d_{1,j'=2}}{d_{1,j=2} + d_{1,j'=2} + d_2} + \sum_c \Pr_{(x',y') \in \Psi_k(x,y)}(x' = c2) \frac{1}{d_{1,j'=2}} \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \frac{d_{1,j'=2}}{d_{1,j=2}(1 + d_{1,j'=2})} + \frac{d_{1,j=2}}{d_{1,j=2} + d_{1,j'=2} + d_2} \frac{1}{d_{1,j'=2}} \right\} \\
&\leq \max_{(x,y) \in S_i} \left\{ \frac{1}{1 + d_{1,j=2}} + \frac{d_{1,j=2}}{d_{1,j=2}(1 + d_{1,j'=2})} \frac{1}{d_{1,j'=2}} \right\} \\
&\leq \frac{1}{q - \Delta + 1} + \frac{1}{(q - \Delta + 1)^2}
\end{aligned}$$

which implies the statement of the lemma. $\square$

This completes the cases of the coupling and we combine the obtained bounds on $\rho_{i,j}^k$ and $\rho_{i,j'}^k$ in the following corollary of Lemmas 14, 15, 16 and 17 which we use in establishing the mixing time of $\mathcal{M}_{\text{edge}}$.

Corollary 18. *Let j and j' be the endpoints of an edge Θ_k . If $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$ then*

$$\rho_{i,j}^k \leq \frac{1}{q-\Delta} + \frac{1}{(q-\Delta)^2} \text{ and } \rho_{i,j'}^k \leq \frac{1}{q-\Delta} + \frac{1}{(q-\Delta)^2}.$$

Remark. Note that the bound in Corollary 18 is never tight. This bound could be improved, however this would only allow us to beat the 2Δ bound for special graphs since the bounds in Lemma 10 are tight.

We are now ready to present a proof of Theorem 3.

Theorem 3. *Let G be a graph with maximum vertex-degree Δ . If $q \geq 2\Delta$ then*

$$\text{Mix}(\mathcal{M}_{\text{edge}}, \varepsilon) \leq \Delta^2 \log(n\varepsilon^{-1}).$$

Proof. Let j and j' be the endpoints of an edge represented by a block Θ_k . Let $\alpha_j = \sum_i \rho_{i,j}^k$ be the influence on site j and $\alpha_{j'} = \sum_i \rho_{i,j'}^k$ then influence on j' . Then $\alpha = \max(\alpha_j, \alpha_{j'})$. Suppose that Θ_k is adjacent to t triangles, that is there are t sites $i_1, \dots, i_t$ such that $\{i, j\} \in E(G)$ and $\{i, j'\} \in E(G)$ for each $i \in \{i_1, \dots, i_t\}$. Note that $0 \leq t \leq \Delta - 1$. There are at most $\Delta - 1 - t$ sites adjacent to j that are not adjacent to j' and at most $\Delta - 1 - t$ sites adjacent to j' that are not adjacent to j . From Lemma 10 a site adjacent only to j will emit an influence of at most $\frac{1}{q-\Delta}$ on site j and Lemma 10 also guarantees that a site only adjacent to j' can emit an influence at most $\frac{1}{(q-\Delta)^2}$ on site j . Corollary 18 says that a site adjacent to both j and j' can emit an influence of at most $\frac{1}{q-\Delta} + \frac{1}{(q-\Delta)^2}$ on site j and hence

$$\begin{aligned} \alpha_j &\leq t \left(\frac{1}{q-\Delta} + \frac{1}{(q-\Delta)^2} \right) + (\Delta - 1 - t) \left(\frac{1}{q-\Delta} \right) + (\Delta - 1 - t) \left(\frac{1}{(q-\Delta)^2} \right) \\ &= \frac{\Delta - 1}{q - \Delta} + \frac{\Delta - 1}{(q - \Delta)^2} \end{aligned}$$

and similarly by considering the influence on site j' we find that

$$\alpha_{j'} \leq \frac{\Delta - 1}{q - \Delta} + \frac{\Delta - 1}{(q - \Delta)^2}.$$

Then using our assumption that $q \geq 2\Delta$ we have

$$\alpha = \max(\alpha_j, \alpha_{j'}) \leq \frac{\Delta - 1}{q - \Delta} + \frac{\Delta - 1}{(q - \Delta)^2} \leq \frac{\Delta - 1}{\Delta} + \frac{\Delta - 1}{\Delta^2} = \frac{\Delta^2 - 1}{\Delta^2} = 1 - \frac{1}{\Delta^2} < 1$$

and we obtain the stated bound on the mixing time by applying Theorem 2. $\square$

4 Application: Colouring a Tree

This section contains the proof of Theorem 4 which improves the least number of colours required for mixing of systematic scan on a tree for individual values of Δ . Recall the definition of the systematic scan $\mathcal{M}_{\text{tree}}$ where the set of blocks Θ is defined as follows. Let the block Θ_k contain a site r along with all sites below r in the tree that are at most $h - 1$ edges away from r . We call

h the *height* of the blocks and h is defined for each Δ in Table 1. The set of blocks Θ covers the sites of the tree and we construct Θ such that no block has height less than h . $P^{[k]}$ is the transition matrix for performing a heat-bath move on block Θ_k and hence $P^{[k]}(x)$ is the uniform distribution on the set of configurations that agree with x off Θ_k and where no edge incident to a site in Θ_k is monochromatic. The transition matrix of the Markov chain $\mathcal{M}_{\text{tree}}$ is $\prod_{k=1}^m P^{[k]}$ where m is the number of blocks.

We will use standard terminology when discussing the structure of the tree. In particular will say that a site i is a *descendant* of a site j (or j is a *predecessor* of i) if j is on the simple path from the root of the tree to i . We will call a site j a *child* of a site i (or i is the *parent* of j) if i and j are adjacent and j is a descendant of i . Finally $N_k(j) = \{i \in \partial\Theta_k \mid i \text{ is a descendant of } j\}$ is the set of descendants of j on the boundary of Θ_k .

Let $(x, y) \in S_i$ where i is on the boundary of some block Θ_k . The following lemma will provide upper bounds on the probability of disagreement at any site in the block.

Lemma 19. *Let $(x, y) \in S_i$ and suppose that i is adjacent to exactly one site in a block Θ_k . Then there exists a coupling ψ of $D_1 = P^{[k]}(x)$ and $D_2 = P^{[k]}(y)$ in which*


$$\Pr_{(x', y') \in \psi}(x'_j \neq y'_j) \leq \frac{1}{(q - \Delta)^{d(i, j)}}$$

for all $j \in \Theta_k$ where $d(i, j)$ is the edge distance from i to j .

Proof. We construct a coupling ψ of D_1 and D_2 based on the recursive coupling defined in Goldberg et al. [10]. The following definitions are based on Figure 12. Let $R \subseteq V$ be a set of sites. Also let (X, X') be a pair of colourings of the sites on the boundary of R (recall that the boundary of R is the set of sites that are not included in R but are adjacent to some site in R) which use the same colour for every site, *except* for one site u which is coloured l in X and l' in X' . We then say that $A(R, (X, X'), u, (l, l'))$ is a boundary pair. For a boundary pair $A(R, (X, X'), u, (l, l'))$ we let $v \in R$ be the site in R that is adjacent to u . We think of v as the root of R and note that we may need to turn the original tree “upside down” in order to achieve this, however the meaning should be clear. We then label the children (in R) of v as $v_1, \dots, v_d$ and let $T = \{R_1, \dots, R_d\}$ be set the of d subtrees of R that do not contain site v , that is for $R_k \in T$ we define $R_k = \{j \in R \mid j = v_k \text{ or } j \text{ is a descendant of } v_k\}$. Finally let D and D' be the uniform distributions on colourings of R consistent with the boundary colourings X and X' respectively and let $D(v)$ (respectively $D'(v)$) be the uniform distribution on the color at site v induced by D (respectively D'). Then Ψ_R is the recursive coupling of D and D' summarised as follows.

1. If $l = l'$ then the distributions D and D' are the same and we use the identity coupling, in which the same colouring is used in both copies. Otherwise we couple $D(v)$ and $D'(v)$ greedily to maximise the probability of assigning the same colour to site v in both distributions. If R consists of a single site then this completes the coupling.
2. Suppose that the pair of colours (c, c') were drawn for v in the coupling from step 1. For each subtree $R' \in \{R_1, \dots, R_d\}$ we have a well defined boundary pair $A(R', (X_{R'}, X'_{R'}), v, (c, c'))$ where $X_{R'}$ is the boundary colouring X restricted to the sites on the boundary of R' . For each pair of colours (c, c') and $R' \in T$ we recursively construct a coupling $\Psi_{R'}(c, c')$ of the distributions induced by the boundary pair $A(R', (X_{R'}, X'_{R'}), v, (c, c'))$.

Figure 12: The region defined in a boundary pair and the construction of the subtrees.



Initially we let the boundary pair be $A(R = \Theta_k, (X = x, Y = y), u = i, (l = x_i, l' = y_i))$ and our coupling ψ of D_1 and D_2 is thus the recursive coupling Ψ_{Θ_k} constructed above.

We prove the statement of the lemma by induction on $d(i, j)$. The base case is $d(i, j) = 1$. Applying Lemma 13 from Goldberg et al. [10] we can upper bound the probability of $x'_j \neq y'_j$ where (x', y') is drawn from ψ by assigning the worst possible colouring to neighbours of j in Θ_k . Site j has at most $\Delta - 1$ neighbours (other than i) so there are at least $q - \Delta$ colours available for j in both distributions. There is also at most one colour which is valid for j in x but not in y (and vice versa) so

$$\Pr_{(x', y') \in \psi}(x'_j \neq y'_j) \leq \frac{1}{q - \Delta}.$$

Now let R' be the subtree of Θ_k containing site j and let v be the site in Θ_k adjacent to i . Assume that for $d(v, j) = d(i, j) - 1$

$$\Pr_{(x', y') \in \Psi_{R'}(c, c')}(x'_j \neq y'_j) \leq \frac{1}{(q - \Delta)^{d(v, j)}}.$$

Now for $(x, y) \in S_i$

$$\begin{aligned}
\Pr_{(x', y') \in \psi}(x'_j \neq y'_j) &= \Pr_{(x', y') \in \Psi_{\Theta_k}}(x'_j \neq y'_j) \\
&= \sum_{\substack{c, c' \\ c \neq c'}} \Pr_{(x', y') \in \Psi_R}(x'_v = c, y'_v = c') \Pr_{(x', y') \in \Psi_{R'}(c, c')}(x'_j \neq y'_j) \\
&\leq \frac{1}{(q - \Delta)^{d(i, j) - 1}} \sum_{\substack{c, c' \\ c \neq c'}} \Pr_{(x', y') \in \Psi_R}(x'_v = c, y'_v = c') \\
&\leq \frac{1}{(q - \Delta)^{d(i, j)}}
\end{aligned}$$

where the first inequality is the inductive hypothesis and the last is a consequence of the base case. $\square$

We will now use the coupling from Lemma 19 to define the coupling $\Psi_k(x, y)$ of the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$ for $(x, y) \in S_i$. If $i \in \partial\Theta_k$ then it is adjacent to exactly one site in Θ_k and we use the coupling from Lemma 19. If $i \notin \partial\Theta_k$ then the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$ are the same since we are using heat-bath updates and so we can use the identity coupling. We summarise the bounds on $\rho_{i, j}^k$ in the following corollary of Lemma 19, which we will use in the proof of Theorem 4.

Corollary 20. *Let $d(i, j)$ denote the number of edges between i and j . Then for $j \in \Theta_k$*

$$\rho_{i, j}^k \leq \begin{cases} \frac{1}{(q - \Delta)^{d(i, j)}} & \text{if } i \in \partial\Theta_k \\ 0 & \text{otherwise.} \end{cases}$$

Theorem 4. *Let G be a tree with maximum vertex-degree Δ . If $q \geq f(\Delta)$ where $f(\Delta)$ is specified in Table 1 for small Δ then*

$$\text{Mix}(\mathcal{M}_{\text{tree}}, \varepsilon) = O(\log(n\varepsilon^{-1})).$$

Proof. We will use Theorem 2 and assign a weight to each site i such that $w_i = \xi^{d_i}$ where d_i is the edge distance from i to the root and ξ is defined in Table 1 for each Δ . For a block Θ_k and $j \in \Theta_k$ we let


$$\alpha_{k, j} = \frac{\sum_i w_i \rho_{i, j}^k}{w_j}$$

denote the total weighted influence on site j when updating block Θ_k . For each block Θ_k and each site $j \in \Theta_k$ we will upper bound $\alpha_{k, j}$ and hence obtain an upper bound on $\alpha = \max_k \max_{j \in \Theta_k} \alpha_{k, j}$. Note from Corollary 20 that $\rho_{i, j}^k = 0$ when $i \in \Theta_k$ so we only need to bound $\rho_{i, j}^k$ for $i \in \partial\Theta_k$.

We first consider a block Θ_k that does not contain the root. The following labels refer to Figure 13 in which a solid line is an edge and a dotted line denotes the existence of a simple path between two sites. Let $p \in \partial\Theta_k$ be the predecessor of all sites in Θ_k and $d_r - 1$ be the distance from p to the root of the tree i.e., $w_p = \xi^{d_r - 1}$. The site $r \in \Theta_k$ is a child of p . Now consider a site $j \in \Theta_k$ which has distance d to r , hence $w_j = \xi^{d + d_r}$ and $d(j, p) = d + 1$. From Corollary 20 it then follows that the weighted influence of p on j when updating Θ_k is at most

$$\rho_{p, j}^k \frac{w_p}{w_j} \leq \frac{1}{(q - \Delta)^{d(j, p)}} \frac{\xi^{d_r - 1}}{\xi^{d + d_r}} = \frac{1}{(q - \Delta)^{d + 1}} \frac{1}{\xi^{d + 1}}.$$

Figure 13: A block in the tree. A solid line indicates an edge and a dotted line the existence of a path.



Now consider some site $u \in N_k(j)$ which is on the boundary of Θ_k . Since $u \in N_k(j)$ it has weight $w_u = \xi^{d_r+h}$ and so $d(j, u) = h - d$. Hence Corollary 20 says that the weighted influence of u on j is at most

$$\rho_{u,j}^k \frac{w_u}{w_j} \leq \frac{1}{(q - \Delta)^{d(j,u)}} \frac{\xi^{d_r+h}}{\xi^{d_r+d}} = \frac{1}{(q - \Delta)^{h-d}} \xi^{h-d}.$$

Every site in Θ_k has at most $\Delta - 1$ children so the number of sites in $N_k(j)$ is at most $|N_k(j)| \leq (\Delta - 1)^{h-d}$ and so, summing over all sites $u \in N_k(j)$, the total weighted influence on j from sites in $N_k(j)$ when updating Θ_k is at most

$$\sum_{u \in N_k(j)} \rho_{u,j}^k \frac{w_u}{w_j} \leq \sum_{u \in N_k(j)} \frac{1}{(q - \Delta)^{h-d}} \xi^{h-d} \leq \frac{(\Delta - 1)^{h-d}}{(q - \Delta)^{h-d}} \xi^{h-d}.$$

The influence on j from sites in $\partial\Theta_k \setminus (N_k(j) \cup \{p\})$ will now be considered. These are the sites on the boundary of Θ_k that are neither descendants or predecessors of j . For each site v between j and p , we will bound the influence on site j from sites $b \in N_k(v)$ that contain v on the simple path between b and j . We call this the influence on j via v . Referring to Figure 13 let $v \in \Theta_k$ be a predecessor of j such that $d(j, v) = l$ and observe that v is on level $d_r + d - l$ in the tree and also

that $1 \leq l \leq d$ since v is between p and j in the tree. If v is not the parent of j (that is $l \neq 1$) then let j' be the child of v which is also a predecessor of j , that is j' is on the simple path from v to j . If $l = 1$ we let $j' = j$. Also let v' be any child of v other than j' and observe that v' and j' are both on level $d_r + d - l + 1$. Now let $b \in N_k(v')$ be a descendant of v' and note as before that $w_b = \xi^{d_r+h}$. The distance between b and v' is

$$d(v', b) = d_r + h - (d_r + d - l + 1) = h - d + l - 1$$

and so the number of descendants of v' is at most $|N_k(v')| \leq (\Delta - 1)^{h-d+l-1}$ since each site has at most $\Delta - 1$ children. Site v has at most $\Delta - 2$ children other than j' so the number of sites on the boundary of Θ_k that are descendants of v but not j' is at most

$$|N_k(v) \setminus N_k(j')| \leq (\Delta - 2)|N_k(v')| \leq (\Delta - 2)(\Delta - 1)^{h-d+l-1}.$$

Finally the only simple path from b to j goes via v and the number of edges on this path is

$$d(j, b) = d(j, v) + d(v, v') + d(v', b) = l + 1 + (h - d + l - 1) = h - d + 2l$$

so, using Corollary 20, the weighted influence of b on site j when updating block Θ_k is at most

$$\rho_{b,j}^k \frac{w_b}{w_j} \leq \frac{\xi^{d_r+h}}{\xi^{d_r+d}} \frac{1}{(q - \Delta)^{d(j,b)}} \leq \frac{\xi^{h-d}}{(q - \Delta)^{h-d+2l}}$$

and summing over all descendants of v (other than descendants of j') on the boundary of Θ_k we find that the influence on j via site v is at most

$$\sum_{b \in N_k(v) \setminus N_k(j')} \rho_{b,j}^k \frac{w_b}{w_j} \leq \sum_{b \in N_k(v) \setminus N_k(j')} \frac{\xi^{h-d}}{(q - \Delta)^{h-d+2l}} \leq \xi^{h-d} \frac{(\Delta - 2)(\Delta - 1)^{h-d+l-1}}{(q - \Delta)^{h-d+2l}}. \quad (5)$$

Summing (5) over $1 \leq l \leq d$ gives an upper bound on the the total weighted influence of sites in $\partial\Theta_k \setminus (N_k(j) \cup \{p\})$ on site j when updating Θ_k


$$\sum_{b \in \partial\Theta_k \setminus (N_k(j) \cup \{p\})} \rho_{b,j}^k \frac{w_b}{w_j} \leq \xi^{h-d} \sum_{l=1}^d \frac{(\Delta - 2)(\Delta - 1)^{h-d+l-1}}{(q - \Delta)^{h-d+2l}}$$

and adding the derived influences we find that the influence on site j (on level $d_r + d$) when updating Θ_k is at most

$$\begin{aligned} \alpha_{k,j} &= \frac{\rho_{p,j}^k w_p}{w_j} + \sum_{u \in N_k(j)} \frac{\rho_{u,j}^k w_u}{w_j} + \sum_{b \in \partial\Theta_k \setminus (N_k(j) \cup \{p\})} \frac{\rho_{b,j}^k w_b}{w_j} \\ &\leq \frac{1}{(q - \Delta)^{d+1}} \frac{1}{\xi^{d+1}} + \frac{(\Delta - 1)^{h-d}}{(q - \Delta)^{h-d}} \xi^{h-d} + \xi^{h-d} \sum_{l=1}^d \frac{(\Delta - 2)(\Delta - 1)^{h-d+l-1}}{(q - \Delta)^{h-d+2l}}. \end{aligned}$$

Now consider the block containing the root of the tree, r . Let this be block Θ_0 and note that $w_r = 1$. The only difference between Θ_0 and any other block is that r may have Δ children. There

Figure 14: The influence on site j via the root. A line denotes an edge and a dotted line the existence of a simple path.



are at most $\Delta(\Delta - 1)^{h-1}$ descendants of r in $\partial\Theta_0$, each of which has weight ξ^h so, using Corollary 20, the weighted influence on the root is at most

$$\alpha_{0,r} = \sum_{b \in N_0(r)} \rho_{b,r}^0 \frac{w_b}{w_r} \leq \frac{\Delta(\Delta - 1)^{h-1}}{(q - \Delta)^h} \xi^h.$$

Now consider a site j on level $d \neq 0$ in block Θ_0 . As in the general case considered above there is an influence of at most

$$\sum_{b \in N_0(j)} \frac{\rho_{b,j}^0 w_b}{w_j} \leq \frac{(\Delta - 1)^{h-d}}{(q - \Delta)^{h-d}} \xi^{h-d}$$

on j from the sites in $N_0(j)$. Now consider the influence on site j from $\partial\Theta_0 \setminus N_0(j)$. We first consider the influence on j via r , which is shown in Figure 14. Site r has at most $\Delta - 1$ children other than the site j' which is the child of r that is on the path from r to j . Each child of r has at most $(\Delta - 1)^{h-1}$ descendants in $\partial\Theta_0$ and each such descendant has distance $h + d$ to j . Hence, from Corollary 20, the influence on j via the root is at most

$$\sum_{b \in N_0(r) \setminus N_0(j')} \frac{\rho_{b,j}^0 w_b}{w_j} \leq \sum_{b \in N_0(r) \setminus N_0(j')} \frac{\xi^h}{\xi^d} \frac{1}{(q - \Delta)^{d(b,j)}} \leq \frac{(\Delta - 1)^h}{(q - \Delta)^{h+d}} \xi^{h-d}.$$

Finally consider then influence on j from the remaining sites, which are in the set $R = \partial\Theta_0 \setminus (N_0(j) \cup (N_0(r) \setminus N_0(j')))$. Again consider a site $v \neq r \in \Theta_0$ where v is a predecessor of j and $d(j, v) = l$. In this case we have $1 \leq l \leq d-1$ since $l = d$ is the root which has already been considered. This is the same situation as arose in the general case considered above (see Figure 13) so (5) is an upper bound on the influence on j via v and so summing (5) over $1 \leq l \leq d-1$ and adding the other influences on j we obtain an upper bound on the total weighted influence on site j when updating block Θ_0

$$\begin{aligned} \alpha_{0,j} &= \sum_{b \in N_0(j)} \frac{\rho_{b,j}^0 w_b}{w_j} + \sum_{b \in N_0(r) \setminus N_0(j')} \frac{\rho_{b,j}^0 w_b}{w_j} + \sum_{b \in R} \frac{\rho_{b,j}^0 w_b}{w_j} \\ &\leq \frac{(\Delta-1)^{h-d}}{(q-\Delta)^{h-d}} \xi^{h-d} + \frac{(\Delta-1)^h}{(q-\Delta)^{h+d}} \xi^{h-d} + \xi^{h-d} \sum_{l=1}^{d-1} \frac{(\Delta-2)(\Delta-1)^{h-d+l-1}}{(q-\Delta)^{h-d+2l}}. \end{aligned}$$

We require $\alpha < 1$ which we obtain by satisfying the system of inequalities given by setting

$$\alpha_{k,j} < 1 \tag{6}$$

for all blocks Θ_k and sites $j \in \Theta_k$. In particular we need to find an assignment to ξ and h that satisfies (6) given Δ and q . Table 1 shows the least number of colours $f(\Delta)$ required for mixing for small Δ along with a weight, ξ , that satisfies the system of equations and the required height of the blocks, h . These values were verified by checking the resulting $2h$ inequalities for each Δ using Mathematica. The least number of colours required for mixing in the single site setting is also included in the table for comparison. $\square$

5 A Comparison of Influence Parameters

We conclude with a discussion of our choice of influence parameter α denoting the maximum influence on any site in the graph. As we will be comparing the condition $\alpha < 1$ to the corresponding, but unweighted, conditions in Dyer et al. [5] and Weitz [17] we will let $w_i = 1$ for each site. Recall our definitions (letting $w_i = 1$) of $\rho_{i,j}^k$ and α

$$\rho_{i,j}^k = \max_{(x,y) \in S_i} \{\Pr_{(x',y') \in \Psi_k(x,y)}(x'_j \neq y'_j)\} \text{ and } \alpha = \max_k \max_{j \in \Theta_k} \sum_{i \in V} \rho_{i,j}^k$$

where $\Psi_k(x, y)$ is a coupling of the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$. We have previously stated that this is not the standard way to define the influence of i on j since the coupling is directly included in the definition of ρ . It is worth pointing out, however, that the corresponding definition in Weitz [17], which is also for block dynamics, also makes explicit use of the coupling. In the single site setting (Dyer et al. [5]) the influence of i on j , which we will denote $\hat{\rho}_{i,j}$, is defined by

$$\hat{\rho}_{i,j} = \max_{(x,y)_i} d_{\text{TV}}(\mu_j(x), \mu_j(y))$$

where $\mu_j(x)$ is the distribution on spins at site j induced by $P^{[j]}(x)$. The corresponding condition is $\hat{\alpha} = \max_j \sum_{i \in V} \hat{\rho}_{i,j} < 1$. We will show (Lemma 21) that $\hat{\rho}_{i,j}$ is a special case of $\rho_{i,j}^j$ when $\Theta_j = \{j\}$ and $\Psi_j(x, y)$ is a coupling minimising the Hamming distance at site j . This will prove our claim

that our condition $\alpha < 1$ is a generalisation of the single site condition $\hat{\alpha} < 1$. Before establishing this claim we discuss the need to include the coupling explicitly when working with block dynamics. Consider a pair of distinct sites $j \in \Theta_k$ and $j' \in \Theta_k$ and a pair of configurations $(x, y) \in S_i$. When updating block Θ_k the dynamics needs to draw a pair of new configurations (x', y') from the distributions $P^{[k]}(x)$ and $P^{[k]}(y)$ as previously specified. Hence the interaction between j and j' has to be according to these distributions and so it is not possible to consider the influence of i on j and the influence of i on j' separately. In the context of our definition of ρ this means that the influence of i on j and the influence of i on j' have to be defined using the *same coupling*. This is to say that the coupling $\Psi_k(x, y)$ can only depend on the block Θ_k and the initial pair of configurations x and y , which in turn specify which site is labeled i . It is important to note that the coupling can not depend on j , since otherwise having a small influence on a site would not imply rapid mixing of systematic scan (or indeed random update). The reason why we need to make this distinction when working with block dynamics but not the single site dynamics is that in the single site setting $\hat{\rho}_{i,j}$ is the influence of site i on j when updating site j and hence whichever coupling is used must implicitly depend on j . Since the coupling can depend on j in the single site case it is natural to use the “optimal” coupling, which minimises the probability of having a discrepancy at site j . By definition of total variation distance, the probability of having a discrepancy at site j under the optimal coupling is $d_{\text{TV}}(\mu_j(x), \mu_j(y)) = \hat{\rho}_{i,j}$ (see e.g. Aldous [1]). We will now show that $\hat{\rho}_{i,j}$ is a special case of $\rho_{i,j}^j$ in the way described above.

Lemma 21. *Suppose that for each site $j \in V$ we have a block $\Theta_j = \{j\}$ and that $\Theta = \{\Theta_j\}_{j=1,\dots,n}$. Also suppose that for each pair $(x, y) \in S_i$ of configurations $\Psi_j(x, y)$ is a coupling of $P^{[j]}(x)$ and $P^{[j]}(y)$ in which, for each $c \in C$,*

$$\Pr_{(x', y') \in \Psi_j(x, y)}(x'_j = y'_j = c) = \min(\Pr_{\mu_j(x)}(c), \Pr_{\mu_j(y)}(c))$$

where $\Pr_{\mu_j(x)}(c)$ is the probability of drawing colour c from distribution $\mu_j(x)$. Then $\rho_{i,j}^j = \hat{\rho}_{i,j}$.

Proof. To see that $\Psi_j(x, y)$ is a coupling of $P^{[j]}(x)$ and $P^{[j]}(y)$ it is sufficient to observe that $\Pr_{x' \in P^{[j]}(x)}(x'_j = c) = \Pr_{\mu_j(x)}(c)$ and similarly $\Pr_{y' \in P^{[j]}(y)}(y'_j = c) = \Pr_{\mu_j(y)}(c)$ since j is the only

site in Θ_j . Thus we have

$$\begin{aligned}
\rho_{i,j}^j &= \max_{(x,y) \in S_i} \left\{ \Pr_{(x',y') \in \Psi_j(x,y)}(x'_j \neq y'_j) \right\} \\
&= \max_{(x,y) \in S_i} \left\{ 1 - \sum_{c \in C} (\Pr_{(x',y') \in \Psi_j(x,y)}(x'_j = y'_j = c)) \right\} \\
&= \max_{(x,y) \in S_i} \left\{ 1 - \sum_{c \in C} \min(\Pr_{\mu_j(x)}(c), \Pr_{\mu_j(y)}(c)) \right\} \\
&= \max_{(x,y) \in S_i} \left\{ \sum_{c \in C} \Pr_{\mu_j(x)}(c) - \min(\Pr_{\mu_j(x)}(c), \Pr_{\mu_j(y)}(c)) \right\} \\
&= \max_{(x,y) \in S_i} \left\{ \sum_{c \in C^+} \Pr_{\mu_j(x)}(c) - \Pr_{\mu_j(y)}(c) \right\} \\
&= \max_{(x,y) \in S_i} \left\{ \frac{1}{2} \sum_{c \in C} |\Pr_{\mu_j(x)}(c) - \Pr_{\mu_j(y)}(c)| \right\} \\
&= \max_{(x,y) \in S_i} d_{\text{TV}}(\mu_j(x), \mu_j(y)) \\
&= \hat{\rho}_{i,j}
\end{aligned}$$

where $C^+ = \{c \mid \Pr_{\mu_j(x)}(c) \geq \Pr_{\mu_j(y)}(c)\}$. □

Finally we will show that the condition corresponding to $\alpha < 1$ in Weitz's paper [17] does not imply rapid mixing of systematic scan. Let $B(j)$ be the set of block indices that contain site j and $b(j)$ the size of this set. Weitz refers to the sum $\sum_{k \in B(j)} \sum_i \rho_{i,j}^k$ as the total influence *on* site j and the parameter representing the maximum influence on a site, which we denote α_W to distinguish it from our own definition of α , is defined as

$$\alpha_W = \max_j \sum_{k \in B(j)} \sum_i \frac{\rho_{i,j}^k}{b(j)}.$$

We note that the single site influence parameter $\hat{\alpha}$ used in Dyer et al. [5] to prove rapid mixing of systematic scan is a special case of α_W when the coupling from Lemma 21 is used and each site is contained in exactly one block of size one.

It is proved in Weitz [17] that the condition $\alpha_W < 1$ implies spatial mixing of a random update Markov chain and hence that the Gibbs measure is unique. We will now show that the parameters α and α_W are different and in particular that the condition $\alpha_W < 1$ does not imply rapid mixing of systematic scan. To show this we exhibit a spin system for which a systematic scan Markov chain does not mix rapidly but $\alpha_W < 1$. It is sufficient to show that a specific systematic scan does not mix, since Theorem 2 states that *any* systematic scan with a specified set of blocks mixes when $\alpha < 1$.

Observation 22. *There exists a spin system for which $\alpha_W < 1$ and $\alpha = 1$ but systematic scan does not mix.*

Consider the following spin system. Let G be the n -vertex cycle and label the sites $0, \dots, n-1$ and C be the set of q spins. Then Θ_i (which has an associated transition matrix $P^{[i]}$) is the block containing site i and $i+1 \pmod n$ and it is updated as follows:

1. The spin at site i is copied to site $i+1$;
2. a spin is assigned to site i uniformly at random from the set of all spins.

The stationary distribution, π , of the spin system is the uniform distribution on all configurations of G . Clearly $P^{[i]}$ satisfies property (1) of the update rule, namely that only sites within the block may change during the update. To see that π is invariant under each $P^{[i]}$ observe that site $i+1$ takes the spin of site i in the original configuration and site j receives a spin drawn uniformly at random. This ensures that each site has probability $1/q$ of having each spin and that they are independent.

We define the ρ values for this spin system by using the following coupling. Consider a block Θ_j for update. The spin at site $j+1$ is deterministic in both copies, and each copy selects the same colour for site j when drawing uniformly at random from C . First suppose that site j is the discrepancy between two configurations. Then, since the spin at j is copied to site $j+1$, the spin of site $j+1$ becomes a disagreement in the coupling and hence $\rho_{j,j+1}^j = 1$. The spin at j is drawn uniformly at random from C in both copies and coupled perfectly so $\rho_{j,j}^j = 0$. Now suppose that the two configurations differ at a site $i \neq j$. Then $\rho_{i,j+1}^j = 0$ since both configurations have the same colour for site j , and $\rho_{i,j}^j = 0$ since the spins at site j are coupled perfectly. Using the values of ρ we deduce that

$$\alpha_W = \max_j \sum_{k \in B(j)} \sum_i \frac{\rho_{i,j}^k}{b(j)} = \frac{1}{2} \left(\rho_{j-1,j}^{j-1} + \sum_{i \neq j-1} \rho_{i,j}^{j-1} + \sum_i \rho_{i,j}^j \right) = \frac{1}{2}$$

and $\alpha = \max_k \max_{j \in \Theta_k} \sum_i \rho_{i,j}^k = 1$.

Let $\mathcal{M}_\rightarrow$ be the systematic scan Markov chain that updates the blocks in the order $\Theta_0, \Theta_1, \dots, \Theta_{n-1}$. For each block Θ_i note that if a configuration y is obtained from updating block Θ_i starting from x then $y_{i+1} = x_i$. Hence when performing the systematic scan, the spin of site 0 in the original configuration moves around the ring ending at site $n-1$ before the update of block Θ_{n-1} moves it on to site 0. Hence if configuration x' is obtained from one complete scan starting from a configuration x we have $x'_0 = x_0$ and the systematic scan Markov chain does not mix since site 0 will always be assigned the same spin after each complete scan.

Acknowledgments

I am grateful to Leslie Goldberg for several useful discussions regarding technical issues and for providing detailed and helpful comments on a draft of this article. I would also like to thank Paul Goldberg for useful comments during the early stages of this work.

References

- [1] David J Aldous. Random walks on finite groups and rapidly mixing markov chains. In *Séminaire de probabilités XVII*, pages 243–297. Springer-Verlag, 1983.

- [2] Magnus Bordewich, Martin Dyer, and Marek Karpinski. Stopping times, metrics and approximate counting. In Michele Bugliesi, Bart Preneel, Vladimiro Sassone, and Ingo Wegener, editors, *ICALP*, volume 4051 of *Lecture Notes in Computer Science*, pages 108–119. Springer, 2006.
- [3] Roland Lvovich Dobrushin. Prescribing a system of random variables by conditional distributions. *Theory Prob. and its Appl.*, 15:458–486, 1970.
- [4] Roland Lvovich Dobrushin and Senya B. Shlosman. Constructive criterion for the uniqueness of Gibbs field. In Jozsef Fritz, Arthur Jaffe, and Domokos Szasz, editors, *Statistical mechanics and dynamical systems*, volume 10 of *Progress in Physics*, pages 371–403. Birkhäuser, Boston, 1985.
- [5] Martin Dyer, Leslie Ann Goldberg, and Mark Jerrum. Dobrushin conditions and systematic scan. In Josep Díaz, Klaus Jansen, José D. P. Rolim, and Uri Zwick, editors, *APPROX-RANDOM*, volume 4110 of *Lecture Notes in Computer Science*, pages 327–338. Springer, 2006.
- [6] Martin Dyer, Leslie Ann Goldberg, and Mark Jerrum. Matrix norms and rapid mixing for spin systems. *ArXiv math.PR/0702744*, 2006.
- [7] Martin Dyer, Leslie Ann Goldberg, and Mark Jerrum. Systematic scan and sampling colourings. *Annals of Applied Probability*, 16(1):185–230, 2006.
- [8] Martin Dyer, Alistair Sinclair, Eric Vigoda, and Dror Weitz. Mixing in time and space for lattice spin systems: A combinatorial view. *Random Structures and Algorithms*, 24(4):461–479, 2004.
- [9] Hans Föllmer. A covariance estimate for Gibbs measures. *J. Funct. Anal.*, 46:387–395, 1982.
- [10] Leslie Ann Goldberg, Russ Martin, and Mike Paterson. Strong spatial mixing for lattice graphs with fewer colours. *SICOMP*, 35(2):486–517, 2005.
- [11] Thomas P. Hayes. A simple condition implying rapid mixing of single-site dynamics on spin systems. In *FOCS*, pages 39–46. IEEE Computer Society, 2006.
- [12] Claire Kenyon, Elchanan Mossel, and Yuval Peres. Glauber dynamics on trees and hyperbolic graphs. In *Proc. 42nd Annual IEEE Symposium on Foundations of Computer Science*, pages 568–578, 2001.
- [13] Fabio Martinelli, Alistair Sinclair, and Dror Weitz. Glauber dynamics on trees: Boundary conditions and mixing time. *Communications in Mathematical Physics*, 250(2):301–334, 2004.
- [14] Barry Simon. *The Statistical Mechanics of Lattice Gases*. Princeton University Press, 1993.
- [15] Eric Vigoda. Improved bounds for sampling colourings. *J. Math. Phys.*, 2000.
- [16] Dror Weitz. *Mixing in Time and Space for Discrete Spin Systems*. PhD thesis, University of California, Berkley, 2004.
- [17] Dror Weitz. Combinatorial criteria for uniqueness of Gibbs measures. *Random Structures and Algorithms*, 27(4):445–475, 2005.